

Table 3: Comparison of low versus high numbers of infiltrating immune cells in OSCC patients* with low and high PD-L1 scoring using Fisher's exact test.

	PD-L1 low <i>no. (%)</i>	PD-L1 high <i>no. (%)</i>	<i>P</i>
CD3			
Low	10 (37.0)	4 (22.2)	0.237
High	17 (63.0)	14 (77.8)	
CD4			
Low	21 (80.8)	8 (47.1)	0.025
High	5 (19.2)	9 (52.9)	
CD8			
Low	18 (69.2)	7 (41.2)	0.066
High	8 (30.8)	10 (58.8)	
CD20			
Low	11 (40.7)	9 (50.0)	0.379
High	16 (59.3)	9 (50.0)	
CD68			
Low	15 (55.6)	9 (50.0)	0.475
High	12 (44.4)	9 (50.0)	
DC-lamp			
Low	14 (58.3)	9 (50.0)	0.411
High	10 (41.7)	9 (50.0)	
Eosinophils			
Low	17 (63.0)	8 (44.4)	0.179
High	10 (37.0)	10 (55.6)	
Mast cells			
Low	13 (48.1)	8 (44.4)	0.525
High	14 (51.9)	10 (55.6)	

* Analyses for CD4+ and CD8+ cells were carried out in 43 patients and for DC-lamp+ cells in 42 patients. All other analyses included 45 patient