

# Greater topoclimatic control of above- versus below-ground communities

Heidi K. Mod<sup>1,2</sup>  | Daniel Scherrer<sup>1,3</sup>  | Valeria Di Cola<sup>1</sup>  | Olivier Broennimann<sup>1,4</sup>  |  
 Quentin Blandenier<sup>5,6</sup>  | Frank T. Breiner<sup>1</sup> | Aline Buri<sup>4</sup> | Jérôme Goudet<sup>1,7</sup>  |  
 Nicolas Guex<sup>8,9</sup>  | Enrique Lara<sup>6</sup>  | Edward A. D. Mitchell<sup>5,10</sup>  |  
 Hélène Niculita-Hirzel<sup>11</sup>  | Marco Pagni<sup>9</sup> | Loïc Pellissier<sup>3,12</sup> | Eric Pinto-Figueroa<sup>13</sup> |  
 Ian R. Sanders<sup>1</sup>  | Benedikt R. Schmidt<sup>14,15</sup>  | Christophe V. W. Seppey<sup>16</sup> |  
 David Singer<sup>5,17</sup>  | Sylvain Ursenbacher<sup>14,18</sup>  | Erika Yashiro<sup>1,19</sup>  |  
 Jan R. van der Meer<sup>19</sup> | Antoine Guisan<sup>1,4</sup> 

<sup>1</sup>Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland

<sup>2</sup>Department of Geosciences and Geography, University of Helsinki, Helsinki, Finland

<sup>3</sup>Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf, Switzerland

<sup>4</sup>Institute of Earth Surface Dynamics, University of Lausanne, Lausanne, Switzerland

<sup>5</sup>Laboratory of Soil Biodiversity, Institute of Biology, University of Neuchâtel, Neuchâtel, Switzerland

<sup>6</sup>Real Jardín Botánico, CSIC, Madrid, Spain

<sup>7</sup>Swiss Institute of Bioinformatics, University of Lausanne, Lausanne, Switzerland

<sup>8</sup>Bioinformatics Competence Center, University of Lausanne, Lausanne, Switzerland

<sup>9</sup>Vital-IT Group, Swiss Institute of Bioinformatics, Lausanne, Switzerland

<sup>10</sup>Jardin Botanique de Neuchâtel, Neuchâtel, Switzerland

<sup>11</sup>Department of Occupational Health and Environment, Center for Primary Care and Public Health (Unisanté), University of Lausanne, Lausanne, Switzerland

<sup>12</sup>Landscape Ecology, Department of Environmental Systems Science, ETH Zürich, Zürich, Switzerland

<sup>13</sup>Terrabiom, Lausanne, Switzerland

<sup>14</sup>Info Fauna Karch, Neuchâtel, Switzerland

<sup>15</sup>Department of Evolutionary Biology and Environmental Studies, University of Zurich, Zurich, Switzerland

<sup>16</sup>Department of Arctic and Marine Biology, University of Tromsø, Tromsø, Norway

<sup>17</sup>Department of Zoology, Institute of Biosciences, University of São Paulo, São Paulo, Brazil

<sup>18</sup>Department of Environmental Sciences, Section of Conservation Biology, University of Basel, Basel, Switzerland

<sup>19</sup>Department of Fundamental Microbiology, University of Lausanne, Lausanne, Switzerland

## Correspondence

Heidi K. Mod, Department of Geosciences and Geography, P.O. Box 64 (Gustaf Hällströmin katu 2), University of Helsinki, Helsinki, FI-00014, Finland.  
 Email: heidi.mod@helsinki.fi

## Funding information

Norges Forskningsråd, Grant/Award Number: 256132 and 270252; Schweizerischer Nationalfonds zur

## Abstract

Assessing the degree to which climate explains the spatial distributions of different taxonomic and functional groups is essential for anticipating the effects of climate change on ecosystems. Most effort so far has focused on above-ground organisms, which offer only a partial view on the response of biodiversity to environmental gradients. Here including both above- and below-ground organisms, we quantified the degree of topoclimatic control on the occurrence patterns of >1,500 taxa and phylotypes along a

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2020 The Authors. *Global Change Biology* published by John Wiley & Sons Ltd

Förderung der Wissenschaftlichen  
Forschung, Grant/Award Number:  
31003A-1528661, 31003A\_163254,  
315230\_184908 and P2NEP3\_178543;  
Community of Madrid, Grant/Award  
Number: 2017-T1/AMB-5210

c. 3,000 m elevation gradient, by fitting species distribution models. Higher model performances for animals and plants than for soil microbes (fungi, bacteria and protists) suggest that the direct influence of topoclimate is stronger on above-ground species than on below-ground microorganisms. Accordingly, direct climate change effects are predicted to be stronger for above-ground than for below-ground taxa, whereas factors expressing local soil microclimate and geochemistry are likely more important to explain and forecast the occurrence patterns of soil microbiota. Detailed mapping and future scenarios of soil microclimate and microhabitats, together with comparative studies of interacting and ecologically dependent above- and below-ground biota, are thus needed to understand and realistically forecast the future distribution of ecosystems.

#### KEYWORDS

animals, climate change, ecosystems, microorganisms, niche model, plants, species distributions, taxonomic group

## 1 | INTRODUCTION

Climate defines the spatial distribution of a wide range of organisms, and the need to understand this relationship is particularly revived by the current threat of climate change on ecosystems (Illan, Gutierrez, & Wilson, 2010; Lenoir & Svenning, 2015; Miller et al., 2018; Pacifici et al., 2015; Parmesan & Yohe, 2003; Scherrer & Körner, 2011; Steidinger et al., 2019; Woodward & Williams, 1987). In addition to their recognized ecological effects, the prevalent use of average annual, seasonal or monthly air temperature- and precipitation-related factors in many studies is common because they are freely available (e.g. CHELSA; Karger et al., 2017), with existing robust future scenarios (IPCC, 2014). Such macroclimate data are often associated with topographic variables that are easily derived from digital elevation models (DEM; Amatulli et al., 2018) to model topoclimate and to describe topography-related habitat characteristics concerning, for example, the distribution of energy (Wang, Qiu, Wang, Wang, & Liu, 2014) and water (Moeslund et al., 2013).

Topoclimatic factors, albeit not capturing microclimatic variations at very fine spatial resolutions (Boulangeat, Gravel, & Thuiller, 2012; Lembrechts, Lenoir, et al., 2019; Mod, Scherrer, Luoto, & Guisan, 2016; Niittynen, Heikkinen, & Luoto, 2018; Scherrer & Guisan, 2019), are important drivers of species distributions. Yet, how their importance varies among taxonomic groups is largely unknown. Most studies on the effects of topoclimate on the distributional patterns of organisms are strongly biased towards a few taxonomic groups (Table 1; Kharouba, McCune, Thuiller, & Huntley, 2013). For instance, plants and some iconic groups of vertebrates and insects have been extensively studied with varying methodologies ranging from basic observational approaches to complex frameworks (Bateman, VanDerWal, Williams, & Johnson, 2012; Bradie & Leung, 2017; Illan et al., 2010; Lawrence et al., 2014; Miller et al., 2018; Mod & Luoto, 2016; Roberts, Nielsen, & Stenhouse, 2014; Seoane, Bustamante, & Diaz-Delgado, 2004; Staniczenko, Sivasubramaniam, Suttle, & Pearson, 2017), but comparatively fewer attempts have been made to assess the drivers of

**TABLE 1** Number of results for ISI Web of Knowledge query<sup>a</sup> of "(climat\* AND (distribution OR occurrence) AND ([Taxonomic group]))"

Taxonomic group	Number of query results
<i>plant* OR tree* OR vegetation</i>	27,593
<i>bird*</i>	2,829
<i>insect*</i>	2,472
<i>mammal*</i>	2,107
<i>lepidoptera* OR butterfly* OR moth*</i>	1,353
<i>reptil* OR lizard* OR turtle* OR tortoise* OR terrapin* OR snake* OR croco*</i>	1,247
<i>amphibia* OR frog* OR toad* OR salamander* OR newt*</i>	1,212
<i>fungi OR fungus</i>	1,139
<i>bacteria*</i>	896
<i>"micro*organism*" OR "micro*biota"</i>	448
<i>orthoptera* OR grasshopper* OR cricket*</i>	177
<i>bombus OR bumblebee* OR bumble*bee*</i>	78
<i>micro*eukaryot* OR protist*</i>	72

<sup>a</sup>Searched on 11 July 2019.

distributions of soil microorganisms (Bradie & Leung, 2017; Lenoir et al., 2020; Lenoir & Svenning, 2015; Pacifici et al., 2015)—likely due to previous methodological limitations (Riesefeld, Schloss, & Handelsman, 2004) and, for some groups (e.g. soil protists), to a lower sampling effort (Caron, Worden, Countway, Demir, & Heidelberg, 2008; Geisen et al., 2017, 2018; Seppey et al., 2020; Wilkinson, 1998). Existing studies have evidenced the importance of soil characteristics, such as pH, nutrient content and moisture availability, for explaining the diversity, biomass and community structure of microorganisms in soils (Bahram et al., 2018; Bates et al., 2013; Fierer & Jackson, 2006; Serna-Chavez, Fierer, & van Bodegom, 2013; Tedersoo et al., 2014). Furthermore, soil temperature has been shown to be decoupled from

air temperature (e.g. Scherrer & Körner, 2011) and to better explain the fine-scale distributions of plants, especially those with small and close-to-ground growth form (Lembrechts, Lenoir, et al., 2019). Soil microclimate might thus be more important than air macroclimate in explaining the distribution of organisms strongly associated to soil conditions. While a direct quantification of local soil conditions could allow to test their role, an indirect approach is to compare the ability of topoclimatic conditions to explain and predict the distribution of organisms across taxonomic groups with contrasted life histories (Kharouba et al., 2013).

Due to the tight direct and indirect couplings among different taxonomic and functional groups, such as biotic interactions and nutrient cycling (Seibold, Cadotte, Maclvor, Thorn, & Muller, 2018; Wardle et al., 2004), understanding the role of climate control on multiple groups is of crucial importance to more realistically forecast the future of biodiversity and entire ecosystems (Cavicchioli et al., 2019; Guo, Feng, et al., 2018; Hagedorn, Gavazov, & Alexander, 2019; Schleuning et al., 2016). In addition to the taxonomic bias, assessing and comparing the role of climate and warming across groups is hindered by the varying sampling designs, study areas and spatial scales used among the existing studies (Rodríguez-Castañeda, Hof, Jansson, & Harding, 2012). A robust comparative quantification of the role of climatic control on distributions across the various above- and below-ground groups composing ecosystems was thus needed.

Here using comprehensive taxon occurrence data from the same mountainous study area in western Switzerland and a robust modelling, evaluation and prediction framework, we aim to: (a) assess to what extent the commonly used and readily available topoclimatic variables explain alone the distribution patterns of six above-ground taxonomic groups (amphibians, reptiles, grasshoppers, butterflies, bumblebees, vascular plants) and three below-ground ones (soil fungi, bacteria and protists); and (b) make baseline predictions and comparisons about how those taxa may respond to climate change. For this, we fitted climate suitability models with the occurrences of each taxon and six topoclimatic variables and compared models' predictive ability and the taxa's spatial predicted probability of occurrence (PPO) under current and future climatic conditions.

## 2 | MATERIALS AND METHODS

### 2.1 | Study area

The study area is a priority area for transdisciplinary research and biodiversity conservation located in the Western Swiss Alps (46°10'–46°30' N; 6°60'–7°10' E). It covers an area of c. 700 km<sup>2</sup> with elevations ranging from 370 to 3,200 m, resulting in heterogeneous climatic and topographic conditions. The annual mean temperature and precipitation sum vary from 8°C and 1,200 mm at 600 m a.s.l. to –5°C and 2,600 mm at 3,000 m a.s.l., respectively (Bouët, 1985). The distribution of energy, water and debris strongly varies with slope and aspect. Landcover is dominated by alpine grasslands, forests, glaciers and agricultural lands. More detailed descriptions of the

study area are provided in Randin et al. (2006), Dubuis et al. (2011) and at [www.unil.ch/centre-montagne](http://www.unil.ch/centre-montagne).

### 2.2 | Data

The data sets used to build the species distribution models (SDMs) and to project habitat suitability under current and future climatic conditions consist of spatially explicit information on biota and topoclimatic conditions (for detailed description of the data, see Appendix S1). The data on biota comprise site-level presence–absence information about species, genera and/or orders from nine taxonomic groups: amphibians, reptiles, grasshoppers, butterflies, bumblebees, plants, fungi, bacteria and protists (Table 2; Figures S1–S10 in Appendix S1). Above-ground species were visually identified based on their phenotype. Amphibian and reptile data sets resulted from point observations, whereas sampling of plants, insects and soil (for amplicon sequencing of microorganisms) was targeted on non-forested sites selected following a random stratified sampling design considering elevation, slope and aspect. Occurrences of soil bacteria, fungi and protists were obtained by amplicon sequencing of the V5 region of 16S, ITS1 and the V4 of the 18S, respectively, region of the ribosomal RNA small subunit gene. About 38,000 operational taxonomic units (OTUs) were taxonomically identified but as this number exceeded the computational power available (especially for postprocessing of spatial projections), we clustered them at genera and order levels.

The topoclimatic variables used were mean annual temperature, annual temperature range, sum of annual precipitation, topographic position index, slope and potential annual solar radiation (see Figures S11–S20 in Appendix S1). These were derived from climate data and DEM (Federal Office of Topography: [swisstopo.ch](http://swisstopo.ch)) as spatial layers at a resolution of 25 m, with climatic data based on both the current conditions (the most recent climate normal: 1981–2010) and the future scenarios (IPCC's SRES A1b and A2 for the periods 2020–2050, 2045–2075 and 2070–2100; Zubler et al., 2014; [meteo swiss.ch](http://meteo.swiss.ch); and Figure S21 in Appendix S1). As the new Representative Concentration Pathways (RCPs; IPCC, 2014) are not yet available for the study area with the needed resolution, and as we do not aim to make realistic future predictions, but to test climate control among taxonomic groups, we consider these scenarios suitable for this study. For comparison between SRES and RCPs, see Rogelj, Meinshausen, and Knutti (2012).

### 2.3 | Data preparation for analyses

To test the predictive power of the topoclimatic variables, we evaluated the models using a split-sampling approach (10x) assigning 80% of the sites to model calibration and the remaining 20% to model evaluation so that each site was used eight times for calibration and twice for evaluation. The two predicted PPO values per evaluation site were then averaged to make a comparison against the observed occurrences.

**TABLE 2** Data sets of the nine taxonomic groups available and used for modelling in the same study area in the Western Swiss Alps

Taxonomic group	n of taxa (total)	n of modelled taxa	n of sites	Prevalence of taxa (mean $\pm$ SD)	Niche breadth of taxa <sup>a</sup> (mean $\pm$ SD)	Data origin
Amphibians (sp.)	14	5	133	0.40 $\pm$ 0.28	1.71 $\pm$ 0.78	Schmidt and Zumbach (2019)
Reptiles (sp.)	12	12	1,144	0.09 $\pm$ 0.08	1.59 $\pm$ 0.35	Pittet (2017)
Grasshoppers (sp.)	41	21	202	0.25 $\pm$ 0.19	1.98 $\pm$ 0.31	Pradervand et al. (2013)
Butterflies (sp.)	140	78	208	0.22 $\pm$ 0.14	2.14 $\pm$ 0.29	Pellissier et al. (2012)
Bumblebees (sp.)	29	20	202	0.25 $\pm$ 0.16	2.24 $\pm$ 0.21	Pellissier et al. (2013)
Plants (sp.)	795	296	909	0.08 $\pm$ 0.08	2.02 $\pm$ 0.32	Dubuis et al. (2011)
Plants (ge)	288	160	909	0.14 $\pm$ 0.14	2.23 $\pm$ 0.36	Dubuis et al. (2011)
Fungi (ge)	190	92	103	0.34 $\pm$ 0.17	2.98 $\pm$ 0.36	Pinto-Figueroa et al. (2019)
Fungi (or)	36	10	198	0.40 $\pm$ 0.28	3.03 $\pm$ 0.11	Pagni et al. (2013)
Bacteria (ge)	758	346	258	0.38 $\pm$ 0.29	2.92 $\pm$ 0.46	Yashiro et al. (2016)
Bacteria (or)	276	124	258	0.46 $\pm$ 0.29	3.04 $\pm$ 0.48	Yashiro et al. (2016)
Protists (ge)	496	285	220	0.38 $\pm$ 0.26	2.86 $\pm$ 0.43	Seppey et al. (2020)
Protists (or)	161	92	220	0.47 $\pm$ 0.31	2.91 $\pm$ 0.36	Seppey et al. (2020)

Abbreviations: ge, genera; or, orders; sp., species.

<sup>a</sup>Unitless; calculated from variation in principal components of environmental conditions occupied; see Section 2.

Approximately half of the taxa with less than eight presences or absences within the calibration data sets and with less than one presence or absence within the evaluation data sets were removed to allow confident parameter estimation and model evaluation (Table 2). To test if the taxon's commonness or tolerance of varying environmental conditions within the study area are related to the magnitude and direction of predicted future changes, we computed prevalence (as a proportion of presence observations within sites; Figure S22 in Appendix S1) and niche breadth per taxon (Figure S23 in Appendix S1). To obtain an estimate of the niche breadth, we first applied a principal component analysis (PCA; using `dudi.pca` from the R-package `ade4`) to the matrix of the six environmental variables over all pixels of the study area. Niche breadth was then calculated as the sum of the standard deviations of the scores of the first and second PCA axes where the taxon is present (Segurado et al., 2011).

## 2.4 | Modelling

To allow consideration of taxa with extreme (i.e. very low or high) prevalence while still being able to incorporate all six environmental variables without over-parametrizing the models, we used Ensemble of Small Models (ESM; Breiner, Guisan, Bergamini, & Nobis, 2015) with the corresponding functions in the R-package `ecospat` (Di Cola et al., 2017). ESM first fit 'small' models by forming all possible bivariate combinations (i.e. two at a time) of the environmental predictors (here,  $\binom{6}{2} = 15$ ). The bivariate models are then combined to create the final ensemble model. Following the recommendations of Breiner, Nobis, Bergamini, and Guisan (2018), we fitted the bivariate models using the Generalized Boosting Model (GBM), Generalized Linear Model (GLM), Artificial

Neural Network and Classification Tree Analysis. The final ensemble for each taxon was built based on the converged bivariate models weighted based on their model performance as measured by AUC (bivariate models with AUC < 0.5 were discarded from the final ensemble models). The default settings of the functions were used, except that all observations, regardless of taxon prevalence, had the same weight.

## 2.5 | Evaluating the predictive power of topoclimatic variables across taxa

The per taxon models were evaluated with AUC (i.e. the area under the ROC curve), the maximum True Skills Statistic (`maxTSS`) and the maximum Cohen's KAPPA (`maxKAPPA`), which do not require converting the predicted PPO to presence/absence predictions (Guisan, Thuiller, & Zimmermann, 2017), as such thresholding is shown to influence the modelling outcome (Benito, Cayuela, & Albuquerque, 2013; Calabrese, Certain, Kraan, & Dormann, 2014; Liu, Berry, Dawson, & Pearson, 2005). In addition, by using multiple metrics with varying prevalence dependencies (Lawson, Hodgson, Wilson, & Richards, 2014; Lobo, Jiménez-Valverde, & Real, 2008; Somodi, Lepesi, & Botta-Dukat, 2017), we avoided the prevalence of taxa affecting the interpretations of the role of topoclimatic variables. The predictor contributions (sensu variable importance in `biomod2`—R-package; Thuiller, Georges, Engler, & Breiner, 2016) of environmental variables per taxon and per technique were defined as the sum of weights (based on AUC) of bivariate models including the variable of interest divided by the sum of all weights. Variable contributions in the final ensemble models were then further calculated as the weighted means of these contributions with weighting based on the overall AUC of the technique.

## 2.6 | Projections under current and future conditions

PPO of each taxon were obtained across the entire study area (i.e. for all of the 25 m × 25 m pixels defined by topoclimatic data) under current and future environmental conditions based on a full model trained with all data (i.e. outside CV used for evaluation). Some areas under current and future scenarios are not analogous to the environmental conditions present at the sampled sites (Figures S1–S10, S24 and S25 in Appendix S1). This implied that the models needed to extrapolate predictions to the environmental space outside the training and evaluation conditions of the model (Fitzpatrick & Hargrove, 2009). Thus, the changes between current and future predictions for each taxon were also quantified in the subset of the study area with analogous environmental conditions to the sites sampled under current and all future scenarios (i.e. the same mask for all taxa of a group was used to exclude non-analogous areas from the current and all future projections; Figure S25 in Appendix S1). Furthermore, forested areas were masked from the spatial projections of open-habitat insects, plants and soil microorganisms. The same forest mask (from Swiss Map Vector 25 BETA; swisstopo.ch) was used for the current and all future projections, which simulates a scenario where the forest cover in the study area would overall remain constant because the Swiss forest policy aims to prevent forest encroachment of pastures (following the 'lowered agriculture production' socio-economic scenario of Gago-Silva, Ray, & Lehmann, 2017). Other scenarios are, however, plausible too and we acknowledge that incorporating land use changes could affect the predictions (Guo, Lenoir, & Bonebrake, 2018). For example, forest expansion (following the 'liberalization' scenario of Gago-Silva et al., 2017) could further restrict the future distributions of grassland taxa (Tasser & Tappeiner, 2002).

The examination of predictions and the changes therein was based directly on the PPO values under current and future conditions, thus avoiding the problematic use of binarization thresholds (Nenzen & Araujo, 2011). To estimate the expected magnitude of change for each taxon in the study area, we calculated the overlap of the PPO values between current and future projections. For each taxon, the overlap was defined based on the sum of differences between current and future PPO values across all pixels, that is, the Schoener's *D* index (here, derived using the 'modOverlap' function in the fuzzySim R-package; Barbosa, 2015):

$$D(p_X, p_Y) = 1 - \frac{1}{2} \left( \sum_i |p_{X,i} - p_{Y,i}| \right),$$

where  $p_{X,i}$  (resp.  $p_{Y,i}$ ) denotes a taxon's PPO in pixel  $i$  proportional to the sum of taxon's PPOs in all pixels in time  $X$  (resp.  $Y$ ; Barbosa, 2015; Warren, Glor, & Turelli, 2008):

$$p_i = q_i / \sum_1^n q_{n-i},$$

where  $q$  is taxon's PPO and  $n$  is number of pixels. The index varies between 1 (i.e. identical current and future predicted distributions) and 0

(i.e. completely non-overlapping current and future predicted distributions), and thus the lower the overlap, the higher the predicted future change in taxon's distribution.

The spatial variation in the magnitude of community changes within each taxonomic group was assessed by first calculating, for each 25 m × 25 m pixel of the study area, the Euclidian distance between the current and future PPO values of all taxa, using the formula for  $n$ -dimensional space:

$$d(p, q) = \sqrt{\sum_{i=1}^n (p_i - q_i)^2},$$

where  $p_i$  is the PPO of taxon  $i$  in the pixel based on the prediction under current conditions,  $q_i$  is the PPO of taxon  $i$  the pixel based on the prediction under future conditions and  $n$  is the number of taxa in the respective taxonomic group. These Euclidian distances per group per pixel per future scenario were then divided by the highest possible distance per group ( $\sqrt{n}$ ; i.e. the case where for all taxa, PPOs would change from 0 to 1 or from 1 to 0, with resulting distances for all taxa being 1). The values representing the magnitude of community change in each pixel then vary between 0 and 1, where a value of 0 means that the PPO of none of the taxa of the respective taxonomic groups changed between current and future prediction, and a value of 1 means that the PPO of all taxa changed from 1 to 0 or from 0 to 1.


Finally, the direction of change (i.e. estimate of gains vs. losses in suitable habitat) was examined for each taxon and future scenario by calculating the mean and median changes in PPOs across the study area and separately at low, mid and high elevations.

## 2.7 | Comparing ESMs to standard SDMs

The standard ensemble SDMs (with commonly used GBM, GLM, Generalized Additive Models = GAM and Random Forest = RF weighted by AUC within Biomod2-platform; Hao, Elith, Guillera-Arroita, & Lahoz-Monfort, 2019; Thuiller et al., 2016; Thuiller, Lafourcade, Engler, & Araújo, 2009) were fitted for the subsets of species and taxonomic groups that included at least 50 presences and absences in each CV-fold of the training data and at least one presence and absence in each fold of the evaluation data. A selection of model evaluations and analyses of the predicted changes were performed to compare the ESMs and standard SDMs.

## 3 | RESULTS

Across all taxa and techniques, the mean ± standard deviation of the model performance is  $0.747 \pm 0.096$  (measured by AUC; for maxTSS and maxKAPPA see Figure S26 in Appendix S2). The model performance of above-ground species ( $0.823 \pm 0.071$ ) is higher than the model performance of below-ground microorganisms ( $0.700 \pm 0.079$  for genera and  $0.708 \pm 0.086$  for orders; Figure 1a; Figure S26 in Appendix S2). The higher model performances are



**FIGURE 1** The role of topoclimate and climate change in defining current and future taxa distributions. (a) Performance as measured with AUC per taxon of the topoclimatic ensemble models. The model performance of above-ground species is significantly better than the model performance of below-ground microorganisms ( $p < .001$ ; measured with Wilcoxon rank sum test). For other evaluation metrics and per technique (Generalized Linear Model, Generalized Boosting Model, Artificial Neural Network, Classification Tree Analysis), and for plant genera and soil microorganism orders, see Figure S26 in Appendix S2. (b) Spatial overlap in predicted probability of occurrence (PPO) between current and future (A2 2085 scenario) projections within the whole study area. For other scenarios, plant genera, soil microorganism orders and environmentally analogous areas, see Figure S28 in Appendix S2. Boxes in boxplots span the 25th–75th quartile, with median (black bar) and mean (orange point) in the middle. Whiskers span the lowest and highest scores, yet in maximum to  $1.5 \times (75\text{th} - 25\text{th quartile})$ ; outlier scores are indicated by black dots

statistically significant when tested with Mann–Whitney  $U$  test ( $U = 271,431$ ,  $p < .001$  when 432 above-ground species are compared to 723 below-ground genera and  $U = 81,934$ ,  $p < .001$  when compared to 226 below-ground orders) and the probabilities of AUC were found to be higher for above-ground species than for below-ground genus or order are 0.87 and 0.83, respectively, based on the common language effect size (McGraw & Wong, 1992). For all taxonomic groups, the annual mean temperature is the most and solar

radiation the least influential topoclimatic variable (Figure S27 in Appendix S2). The contributions of the other four predictors do not rank consistently among the taxonomic groups.


Changes between current and future predicted PPO increase linearly with the severity of the deployed climate scenarios, and the changes are mostly similar between the whole study area and the parts of the study area which exclude novel future environmental conditions (i.e. non-analogous conditions; see Appendix S2). Forecasts show a relatively smaller overlap between the current and future PPO for reptiles, insects and plants, whereas higher overlap indicates less change especially for the soil microbial distributions (Figure 1b; Figure S28 in Appendix S2). The magnitude of overlap is positively correlated with the prevalence and niche breadth of the taxa (see Tables S1 and S2 and Figures S29 and S30 in Appendix S2).

The strongest predicted changes in taxonomic composition across the study area, as measured by the Euclidean distance in the ordination space of the PPO values between the current and future conditions, occur at the highest elevations (Figure 2). This holds for all taxonomic groups, except for amphibians, reptiles and bumblebees, for which the strongest changes are predicted at midelevations (Figure 2d,g,h; Figures S31–S42 in Appendix S2). The PCA, restricted to the non-forested areas, divides the taxonomic groups into five categories (Figure 2j). The future variations in reptile and amphibian compositions strongly differ from the variations of other groups. The change in composition of bumblebees is opposite to that in amphibians. The predicted variation in the compositions of grasshoppers, butterflies, plants and fungi is similar, whereas soil bacteria and protists cluster separately.


There are slightly more taxa with increases than decreases in mean PPO across the study area between now and the future (Figure 3a; Figure S43 in Appendix S2), especially if only the area with analogous environments is considered (Figure 3b; Figure S43 in Appendix S2). For example, under the A2 scenario for 2070–2100, 83.3% of reptile species and 71.7% of bacteria genera showed an increase in mean PPO. In contrast, amphibians, butterflies and plants show the lowest proportional increase in mean PPO (40.0%, 38.5% and 36.5%, respectively). The mean change in PPO across the study area mainly shows a weak and non-significant relationship to the prevalence and niche breadth of taxa (Tables S3 and S4 and Figures S44 and S45 in Appendix S2). Overall, the changes in mean PPO are stronger and often more positive at mid- and high-compared to low elevations (Figure 3c–e; Figure S46 in Appendix S2). Median changes in PPOs are similar to mean changes (Figures S47–S49 in Appendix S3).

For the subset of taxa modelled with both ESMs and standard SDMs (i.e. all variables incorporated simultaneously), the model performances are highly correlated (0.964–0.983; Table S5 and Figure S50 in Appendix S3), but in general, ESMs perform better for most taxonomic groups (Table S6 in Appendix S3). The mean relative variable contributions across taxa follow the same order for ESMs and standard SDMs (Figure S51 in Appendix S3). The mean changes in PPO of ESMs and standard SDMs are highly correlated, but predictions of standard SDMs show a larger variation (Figures S52–S55 and Table S7 in Appendix S3).

**FIGURE 2** Spatial variation in the magnitude of the predicted future changes in community structure. (a) Location of the study area in the Swiss Alps is marked by yellow squares. (b–i, k) Magnitude of predicted changes in community structure between current conditions and future scenario A2 for year 2085. The maximum possible magnitude is 1, meaning that predicted probability of occurrences (PPOs) of all taxa are predicted to change from 1 to 0 or from 0 to 1, and the minimum possible magnitude is 0, meaning that PPOs of none of the taxa are predicted to change between current and future prediction. Grey areas mark the forest cover masked from the predictions of taxa, as sampling targeted non-forested sites. (j) Principal component analysis (PCA) of community structure changes of the nine taxonomic groups in the non-forested areas shows the (dis)similarity in spatial patterns, that is, groups with lines pointing in the same direction have similar spatial patterns in the magnitude of community changes in non-forested areas, whereas lines of varying directions indicate varying patterns. (l) Elevation of the study area. For maps and PCAs of other taxonomic ranks, scenarios and years, see Figures S31–S36 in Appendix S2, and for the relationships between the changes and elevation, and for comparison to analogous environmental space, see Figures S37–S42 in Appendix S2



**FIGURE 3** Proportions of taxa with different mean changes in predicted probability of occurrence (PPO; classified to six classes) under A2 scenario for 2085. (a) Across the total study area. (b) Across an environmentally analogous part of the study area. (c–e) At different elevational bands (low = <1,180 m a.s.l.; mid = 1,180–1,650 m a.s.l.; high > 1,650 m a.s.l.). For other scenarios and taxonomic ranks, see Figures S43 and S46 in Appendix S2. For median changes, see Figures S47–S49 in Appendix S3



## 4 | DISCUSSION

Our results show that topoclimatic variables alone better explain and predict the distributions of above-ground groups (especially reptiles, grasshoppers and butterflies) than those of below-ground microorganisms and that the forecasted future changes in soil microorganism distributions are accordingly smaller and more spatially different. The low model performance indicates that other factors, in addition to the easily obtainable topoclimatic variables, are needed to improve models and future predictions of soil microorganisms (Bahram et al., 2018; Bates et al., 2013; Fierer, Strickland, Liptzin, Bradford, & Cleveland, 2009; Lembrechts, Nijs, & Lenoir, 2019; Li et al., 2018). Importantly, Lembrechts, Lenoir, et al. (2019) showed that soil temperature better explains the distributions of close-to-ground growing plants than free air climate. This suggests that soil microclimate, which better expresses the temperature and moisture conditions affecting below-ground and ground-dwelling organisms and which can be strongly decoupled from air temperature in space and time (see also Bramer et al., 2018; Graae et al., 2012; Scherrer & Körner, 2011; Suggitt et al., 2011), likely also better explains the distributions of soil microorganisms. Furthermore, other edaphic factors (particularly soil pH) and biotic interactions are known to be major drivers for the diversity and community structure of soil fungi and bacteria (Bahram et al., 2018; Bates et al., 2013; Fierer et al., 2009; Li et al., 2018; Malard, Anwar, Jacobsen, & Pearce, 2019; Yashiro et al., 2016), and also but to a lesser extent for the diversity of protists (Seppey et al., 2020). Thus, they can also be expected to strongly contribute to explain the distribution of individual soil microorganisms.

The forecasted lower magnitudes of distributional changes of soil microorganisms, therefore, do not necessarily imply less severe changes to be expected in the future, but a smaller direct role of macroclimatic factors. As the effects of climate change may also be indirect, propagated by alterations, for example, in soil and vegetation (Alexander, Diez, Hart, & Levine, 2016; Collins, Stajich, Weber, Pombubpa, & Diez, 2018; Mod & Luoto, 2016; Yashiro et al., 2018), future scenarios are needed for soil factors in an analogous way to those developed for macroclimate (IPCC, 2014; Lembrechts et al., 2020). Furthermore, the multicollinearity should be assessed between topoclimate and missing predictors (Oliver & Morecroft, 2014; Sears, Raskin, & Angilletta, 2011), because large spatial and temporal offsets have been demonstrated for example between air and soil temperatures (Aalto, Scherrer, Lenoir, Guisan, & Luoto, 2018; Lembrechts, Lenoir, et al., 2019), and thus the used predictors can have a strong influence on the interpretations of future biodiversity changes (Lenoir et al., 2013; Lenoir, Hattab, & Pierre, 2017; Scherrer, Schmid, & Körner, 2011). Deriving and incorporating all influential and change-prone environmental variables at ecologically relevant scales in the models and future predictions (see e.g. Lembrechts, Nijs, et al., 2019; Mod et al., 2016) would allow to assess if the here demonstrated among-group differences in magnitudes (Figure 1b), spatial configurations (Figure 2j) and/or directions (Figure 3) of future distributional changes are realistic. Incongruent

reactivity to environmental changes between biologically dependent taxa and groups (e.g. plants and bacteria; Yashiro et al., 2018) can lead to shuffled species interactions and disruptions in biotic networks (Alexander, Diez, & Levine, 2015; Araújo & Luoto, 2007; Walther, 2010). Furthermore, land use changes, such as expansion of forest cover (Gago-Silva et al., 2017), influence both directly and indirectly (though altering the abiotic conditions) the re-distribution of taxa (Guo, Lenoir, et al., 2018). Thus, in future studies, land use changes should be accounted for together with climate change.

While our analyses revealed important ecological differences in the expected climate change responses among the studied taxonomic groups, three commonalities also emerged. First, among the six topoclimatic variables used, annual mean temperature had the highest average relative contribution in predicting taxa distribution within our study area. While our study area covers a variety of climatic conditions due to the c. 3,000 m elevation range (equivalent to approx. >1,500 km latitudinal gradient; Montgomery, 2006), the generality of this, and our other findings, across ecoregions (Miller et al., 2018; Petitpierre et al., 2016), geographical scales (Birkhofer et al., 2012; Graf, Bollmann, Suter, & Bugmann, 2005) and other groups (e.g. below-ground macrofauna; Decaëns, 2010) remains to be assessed. Second, the taxa with the strongest changes in suitable habitat generally have lower prevalence and narrower environmental niches. However, these changes were not systematically gains or losses of habitat, making these rare specialists sensitive yet not necessarily vulnerable to climate change (Foden et al., 2013; Moritz & Agudo, 2013). Third, we observed the strongest predicted changes in taxa distributions at mid- and high elevations. This pattern has been commonly observed for plants under climate change forecasts in mountain environments (Dirnböck, Dullinger, & Grabherr, 2003; Engler et al., 2011; Steinbauer et al., 2018), and here we show that it applies to other taxonomic groups as well. Nevertheless, this interpretation must be taken with caution because boundary effects are expected to affect the predictions at low elevations. Boundary effects may arise when omitting taxa occurrences from warmer conditions outside the study area (i.e. resulting in erroneous predictions based on truncated response curves; Hannemann, Willis, & Macias-Fauria, 2016) and restricting the taxa to the ones actually found in the study area (i.e. omitting the taxa migrating and invading from lower altitudes as a result of climate tracking; Lenoir, Gégout, Marquet, de Ruffray, & Brisse, 2008; Menéndez, González-Megías, Jay-Robert, & Marquéz-Ferrando, 2014; Petitpierre et al., 2016).

Our study also demonstrated the importance of two methodological developments, which have so far rarely been considered in climate-biodiversity impact studies involving a large number of taxa. Notably, the ESM approach allows more taxa to be included, particularly less frequent and rare taxa, and thus allows more complete modelling of local communities (Breiner et al., 2015, 2018; Lomba et al., 2010), and tends to outperform standard SDMs even for more common species with large sample sizes (Breiner et al., 2015). Interestingly, the predicted changes in PPO were generally lower than those predicted by standard SDMs, a finding never reported before. The likely cause for this is that the bivariate models in ESMs



are less prone to multicollinearity that can bias the estimated effects of variables within standard SDMs, where all predictors are included simultaneously (Graham, 2003). Second, we avoided here the potential bias and loss of information resulting from the traditionally performed thresholding of predictions, because we applied metrics that do not require binarizing probabilities into presences-absences for evaluating the models and assessing the future changes (Jiménez-Valverde & Lobo, 2007; Lawson et al., 2014; Nenzen & Araujo, 2011; Scherrer, Mod, & Guisan, 2020; Vaughan & Ormerod, 2005).

Finally, we would like to point out four insights related to multi-group biodiversity assessments. First, global data sets would be ideal to test the general effects of climate on multiple taxa and groups. However, existing global data sets are strongly biased in terms of taxonomy (Lenoir et al., 2020; Lenoir & Svenning, 2015; Troudet, Grandcolas, Blin, Vignes-Lebbe, & Legendre, 2017) and spatial coverage (Boakes et al., 2010) affecting SDMs and hindering inter-taxa/-group comparisons (Beck, Böller, Erhardt, & Schwanghart, 2014). Here we reduced considerably these potential biases by sampling many taxa through a similar random stratified sampling design (Hirzel & Guisan, 2002) in a same region, where variation in elevation across the study area creates wide environmental gradients. Second, we acknowledge that our models used input data that were taxonomically attributed differently among the studied groups, as species-level information is currently not available and difficult to determine for soil microorganisms unlike for above-ground groups. One notable reason for this is that macroorganisms can be taxonomically catalogued phenotypically, whereas microorganisms were determined from DNA sequencing data only. As a proxy for microorganisms at the species level, we could have used the frequently deployed OTUs, but their total number (>38,000) exceeded the available storage space and computational capacity regarding the post-processing of data (e.g. calculating Euclidian distances in  $n$ -dimensional space). Hence, soil microorganisms were aggregated to the putative level of genera. While this might have added noise in the data for model calibration, we tested this by artificially aggregating the plant species to genera and the microbial genera to orders. This did not systematically affect model performance, and the differences in model performance between taxonomic resolutions (on average 0.06) remained minor compared to the differences in performance between above- and below-ground taxa (on average 0.13). Third, we focused on identifiable groups to present an ecological narrative for key aspects of mountain ecosystems (e.g. amphibians/reptiles—predation, grasshoppers—herbivory, bumblebees—pollination, plants—primary production, fungi—decomposition, bacteria/protists—nutrient cycling), but we acknowledge that interpretations of climate influence might have been different if the taxa had been organized by another grouping (Dormann & Woodin, 2002; Harrison et al., 2010; Peay, Kennedy, & Talbot, 2016; Soudzilovskaia et al., 2013). Fourth, despite using state-of-the-art methodologies, there might be biases in data, models and projections related to taxa detection (Benoit, Jackson, & Ridgway, 2018), sampling methods (e.g. point observations of amphibians and reptiles vs. presence-absence observations of other groups; Aarts, Fieberg, & Matthiopoulos, 2012; field observations

vs. eDNA; Porter & Hajibabaei, 2018; Yoccoz, 2012; including observations also from forested sites for amphibians and reptiles while restricting the analyses to grasslands for other groups) and/or amplification of eDNA (Fernandes, Scherrer, & Guisan, 2019; Pinol, Mir, Gomez-Polo, & Agusti, 2015). However, these issues tend to affect all intergroup biodiversity studies and should not alter specifically our findings compared to all others.

The need to understand the role of climate on biodiversity is particularly invigorated by the current threat of climate change on ecosystems (IPBES, 2019), and realistic future forecasts require incorporating multiple taxonomic and functional groups. Our models indicate both commonalities and differences in climate control among the studied groups. For all taxonomic groups, mean annual temperature was the most important topoclimatic predictor, and the strongest changes were predicted for rare taxa and at high altitudes. However, our results clearly show that the control of topoclimate is stronger on the distribution of above-ground macroorganisms than of below-ground microorganisms, for which soil microclimate likely better expresses temperature and moisture-related constraints. This finding warns against potential disruptions in biotic networks but also implies that we still need to delineate the most ecologically meaningful environmental variables to comprehensively describe the distributions of soil microbiota and how the latter will be affected by climate and soil changes. Due to the important and intricate biological relationships among the different taxonomic groups, realistic forecasts for all members of the ecosystems will be crucial for the optimal assessment of future patterns of biodiversity. For this, future scenarios of other variables than macroclimatic ones, such as soil microclimatic and geochemical conditions, are also needed.

#### ACKNOWLEDGEMENTS

This project was supported by the Swiss National Science Foundation (grant numbers 31003A-1528661 and 315230\_184908 to A.G., P2NEP3\_178543 to D.S. and 31003A\_163254 to E.L. and Q.B.). E.L. was also supported by the programme 'Atracción de talentos' from the Community of Madrid (2017-T1/AMB-5210), and C.V.W.S. acknowledges the support from the Norwegian Research Council projects 270252 (BiodivERsA: Climarctic) and 256132 (ERAnet-LAC: Methanobase).

#### DATA AVAILABILITY STATEMENT














The data that support the findings of this study are available from info fauna karch (for amphibians and reptiles), Dryad (for plants, insects and current environmental conditions at their sampling sites; <https://doi.org/10.5061/dryad.8sf7m0ch5>) and from the authors (soil microorganisms and environmental variables). Restrictions apply to the availability of amphibian and reptile data, which were used under license for this study.

#### ORCID

Heidi K. Mod  <https://orcid.org/0000-0001-7800-2688>

Daniel Scherrer  <https://orcid.org/0000-0001-9983-7510>

Valeria Di Cola  <https://orcid.org/0000-0003-2925-8366>

Olivier Broennimann  <https://orcid.org/0000-0001-9913-3695>  
 Quentin Blandenier  <https://orcid.org/0000-0002-4297-0262>  
 Jérôme Goudet  <https://orcid.org/0000-0002-5318-7601>  
 Nicolas Guex  <https://orcid.org/0000-0001-6023-0519>  
 Enrique Lara  <https://orcid.org/0000-0001-8500-522X>  
 Edward A. D. Mitchell  <https://orcid.org/0000-0003-0358-506X>  
 Hélène Niculita-Hirzel  <https://orcid.org/0000-0003-1467-8819>  
 Ian R. Sanders  <https://orcid.org/0000-0002-9591-8214>  
 Benedikt R. Schmidt  <https://orcid.org/0000-0002-4023-1001>  
 David Singer  <https://orcid.org/0000-0002-4116-033X>  
 Sylvain Ursenbacher  <https://orcid.org/0000-0001-5093-6403>  
 Erika Yashiro  <https://orcid.org/0000-0002-2364-9908>  
 Antoine Guisan  <https://orcid.org/0000-0002-3998-4815>

## REFERENCES

- Aalto, J., Scherrer, D., Lenoir, J., Guisan, A., & Luoto, M. (2018). Biogeophysical controls on soil-atmosphere thermal differences: Implications on warming Arctic ecosystems. *Environmental Research Letters*, 13(7), 074003. <https://doi.org/10.1088/1748-9326/aac83e>
- Aarts, G., Fieberg, J., & Matthiopoulos, J. (2012). Comparative interpretation of count, presence-absence and point methods for species distribution models. *Methods in Ecology and Evolution*, 3(1), 177–187. <https://doi.org/10.1111/j.2041-210X.2011.00141.x>
- Alexander, J. M., Diez, J. M., Hart, S. P., & Levine, J. M. (2016). When climate reshuffles competitors: A call for experimental macroecology. *Trends in Ecology & Evolution*, 31(11), 831–841. <https://doi.org/10.1016/j.tree.2016.08.003>
- Alexander, J. M., Diez, J. M., & Levine, J. M. (2015). Novel competitors shape species' responses to climate change. *Nature*, 525(7570), 515–518. <https://doi.org/10.1038/nature14952>
- Amatulli, G., Domisch, S., Tuanmu, M.-N., Parmentier, B., Ranipeta, A., Malczyk, J., & Jetz, W. (2018). A suite of global, cross-scale topographic variables for environmental and biodiversity modeling. *Scientific Data*, 5, 180040. <https://doi.org/10.1038/sdata.2018.40>
- Araújo, M. B., & Luoto, M. (2007). The importance of biotic interactions for modelling species distributions under climate change. *Global Ecology and Biogeography*, 16(6), 743–753. <https://doi.org/10.1111/j.1466-8238.2007.00359.x>
- Bahram, M., Hildebrand, F., Forslund, S. K., Anderson, J. L., Soudzilovskaia, N. A., Bodegom, P. M., ... Bork, P. (2018). Structure and function of the global topsoil microbiome. *Nature*, 560(7717), 233–237. <https://doi.org/10.1038/s41586-018-0386-6>
- Barbosa, A. M. (2015). fuzzySim: Applying fuzzy logic to binary similarity indices in ecology. *Methods in Ecology and Evolution*, 6(7), 853–858. <https://doi.org/10.1111/2041-210X.12372>
- Bateman, B. L., VanDerWal, J., Williams, S. E., & Johnson, C. N. (2012). Biotic interactions influence the projected distribution of a specialist mammal under climate change. *Diversity and Distributions*, 18(9), 861–872. <https://doi.org/10.1111/j.1472-4642.2012.00922.x>
- Bates, S. T., Clemente, J. C., Flores, G. E., Walters, W. A., Parfrey, L. W., Knight, R., & Fierer, N. (2013). Global biogeography of highly diverse protistan communities in soil. *ISME Journal*, 7(3), 652–659. <https://doi.org/10.1038/ismej.2012.147>
- Beck, J., Böller, M., Erhardt, A., & Schwanghart, W. (2014). Spatial bias in the GBIF database and its effect on modeling species' geographic distributions. *Ecological Informatics*, 19, 10–15. <https://doi.org/10.1016/j.ecoinf.2013.11.002>
- Benito, B. M., Cayuela, L., & Albuquerque, F. S. (2013). The impact of modelling choices in the predictive performance of richness maps derived from species-distribution models: Guidelines to build better diversity models. *Methods in Ecology and Evolution*, 4(4), 327–335. <https://doi.org/10.1111/2041-210X.12022>
- Benoit, D., Jackson, D. A., & Ridgway, M. S. (2018). Assessing the impacts of imperfect detection on estimates of diversity and community structure through multispecies occupancy modeling. *Ecology and Evolution*, 8(9), 4676–4684. <https://doi.org/10.1002/ece3.4023>
- Birkhofer, K., Schöning, I., Alt, F., Herold, N., Klärner, B., Maraun, M., ... Schumpf, M. (2012). General relationships between abiotic soil properties and soil biota across spatial scales and different land-use types. *PLoS One*, 7(8), e43292. <https://doi.org/10.1371/journal.pone.0043292>
- Boakes, E. H., McGowan, P. J. K., Fuller, R. A., Chang-qing, D., Clark, N. E., O'Connor, K., & Mace, G. M. (2010). Distorted views of biodiversity: Spatial and temporal bias in species occurrence data. *PLoS Biology*, 8(6), e1000385. <https://doi.org/10.1371/journal.pbio.1000385>
- Bouët, M. (1985). *Climat et météorologie de la Suisse romande*. Lausanne, Switzerland: Payot.
- Boulangeat, I., Gravel, D., & Thuiller, W. (2012). Accounting for dispersal and biotic interactions to disentangle the drivers of species distributions and their abundances. *Ecology Letters*, 15(6), 584–593. <https://doi.org/10.1111/j.1461-0248.2012.01772.x>
- Bradie, J., & Leung, B. (2017). A quantitative synthesis of the importance of variables used in MaxEnt species distribution models. *Journal of Biogeography*, 44(6), 1344–1361. <https://doi.org/10.1111/jbi.12894>
- Bramer, I., Anderson, B. J., Bennie, J., Bladon, A. J., De Frenne, P., Hemming, D., ... Gillingham, P. K. (2018). Advances in monitoring and modelling climate at ecologically relevant scales. In D. A. Bohan, A. J. Dumbrell, G. Woodward, & M. Jackson (Eds.), *Next generation biomonitoring: Part 1*. Advances in Ecological Research (Vol. 58, pp. 101–161). Cambridge, MA: Academic Press. <https://doi.org/10.1016/bs.aecr.2017.12.005>
- Breiner, F. T., Guisan, A., Bergamini, A., & Nobis, M. P. (2015). Overcoming limitations of modelling rare species by using ensembles of small models. *Methods in Ecology and Evolution*, 6(10), 1210–1218. <https://doi.org/10.1111/2041-210X.12403>
- Breiner, F. T., Nobis, M. P., Bergamini, A., & Guisan, A. (2018). Optimizing ensembles of small models for predicting the distribution of species with few occurrences. *Methods in Ecology and Evolution*, 9(4), 802–808. <https://doi.org/10.1111/2041-210X.12957>
- Calabrese, J. M., Certain, G., Kraan, C., & Dormann, C. F. (2014). Stacking species distribution models and adjusting bias by linking them to macroecological models. *Global Ecology and Biogeography*, 23(1), 99–112. <https://doi.org/10.1111/geb.12102>
- Caron, D. A., Worden, A. Z., Countway, P. D., Demir, E., & Heidelberg, K. B. (2008). Protists are microbes too: A perspective. *The ISME Journal*, 3, 4. <https://doi.org/10.1038/ismej.2008.101>
- Cavicchioli, R., Ripple, W. J., Timmis, K. N., Azam, F., Bakken, L. R., Baylis, M., ... Webster, N. S. (2019). Scientists' warning to humanity: Microorganisms and climate change. *Nature Reviews Microbiology*, 17(9), 569–586. <https://doi.org/10.1038/s41579-019-0222-5>
- Collins, C. G., Stajich, J. E., Weber, S. E., Pombubpa, N., & Diez, J. M. (2018). Shrub range expansion alters diversity and distribution of soil fungal communities across an alpine elevation gradient. *Molecular Ecology*, 27(10), 2461–2476. <https://doi.org/10.1111/mec.14694>
- Decaëns, T. (2010). Macroecological patterns in soil communities. *Global Ecology and Biogeography*, 19(3), 287–302. <https://doi.org/10.1111/j.1466-8238.2009.00517.x>
- Di Cola, V., Broennimann, O., Petitpierre, B., Breiner, F. T., D'Amen, M., Randin, C., ... Guisan, A. (2017). ecospat: An R package to support spatial analyses and modeling of species niches and distributions. *Ecography*, 40(6), 774–787. <https://doi.org/10.1111/ecog.02671>
- Dirnböck, T., Dullinger, S., & Grabherr, G. (2003). A regional impact assessment of climate and land-use change on alpine vegetation. *Journal of Biogeography*, 30(3), 401–417. <https://doi.org/10.1046/j.1365-2699.2003.00839.x>

- Dormann, C. F., & Woodin, S. J. (2002). Climate change in the Arctic: Using plant functional types in a meta-analysis of field experiments. *Functional Ecology*, *16*(1), 4–17. <https://doi.org/10.1046/j.0269-8463.2001.00596.x>
- Dubuis, A., Pottier, J., Rion, V., Pellissier, L., Theurillat, J.-P., & Guisan, A. (2011). Predicting spatial patterns of plant species richness: A comparison of direct macroecological and species stacking modelling approaches. *Diversity and Distributions*, *17*(6), 1122–1131. <https://doi.org/10.1111/j.1472-4642.2011.00792.x>
- Engler, R., Randin, C. F., Thuiller, W., Dullinger, S., Zimmermann, N. E., Araújo, M. B., ... Guisan, A. (2011). 21st century climate change threatens mountain flora unequally across Europe. *Global Change Biology*, *17*(7), 2330–2341. <https://doi.org/10.1111/j.1365-2486.2010.02393.x>
- Fernandes, R. F., Scherrer, D., & Guisan, A. (2019). Effects of simulated observation errors on the performance of species distribution models. *Diversity and Distributions*, *25*(3), 400–413. <https://doi.org/10.1111/ddi.12868>
- Fierer, N., & Jackson, R. B. (2006). The diversity and biogeography of soil bacterial communities. *Proceedings of the National Academy of Sciences of the United States of America*, *103*(3), 626–631. <https://doi.org/10.1073/pnas.0507535103>
- Fierer, N., Strickland, M. S., Liptzin, D., Bradford, M. A., & Cleveland, C. C. (2009). Global patterns in belowground communities. *Ecology Letters*, *12*(11), 1238–1249. <https://doi.org/10.1111/j.1461-0248.2009.01360.x>
- Fitzpatrick, M. C., & Hargrove, W. W. (2009). The projection of species distribution models and the problem of non-analog climate. *Biodiversity and Conservation*, *18*(8), 2255–2261. <https://doi.org/10.1007/s10531-009-9584-8>
- Foden, W. B., Butchart, S. H. M., Stuart, S. N., Vié, J.-C., Akçakaya, H. R., Angulo, A., ... Mace, G. M. (2013). Identifying the world's most climate change vulnerable species: A systematic trait-based assessment of all birds, amphibians and corals. *PLoS One*, *8*(6), e65427. <https://doi.org/10.1371/journal.pone.0065427>
- Gago-Silva, A., Ray, N., & Lehmann, A. (2017). Spatial dynamic modelling of future scenarios of land use change in Vaud and Valais, Western Switzerland. *ISPRS International Journal of Geo-Information*, *6*(4), 115. <https://doi.org/10.3390/ijgi6040115>
- Geisen, S., Mitchell, E. A. D., Adl, S., Bonkowski, M., Dunthorn, M., Ekelund, F., ... Lara, E. (2018). Soil protists: A fertile frontier in soil biology research. *FEMS Microbiology Reviews*, *42*(3), 293–323. <https://doi.org/10.1093/femsre/fuy006>
- Geisen, S., Mitchell, E. A. D., Wilkinson, D. M., Adl, S., Bonkowski, M., Brown, M. W., ... Lara, E. (2017). Soil protistology rebooted: 30 fundamental questions to start with. *Soil Biology & Biochemistry*, *111*, 94–103. <https://doi.org/10.1016/j.soilbio.2017.04.001>
- Graae, B. J., De Frenne, P., Kolb, A., Brunet, J., Chabrierie, O., Verheyen, K., ... Milbau, A. (2012). On the use of weather data in ecological studies along altitudinal and latitudinal gradients. *Oikos*, *121*(1), 3–19. <https://doi.org/10.1111/j.1600-0706.2011.19694.x>
- Graf, R. F., Bollmann, K., Suter, W., & Bugmann, H. (2005). The importance of spatial scale in habitat models: Capercaillie in the Swiss Alps. *Landscape Ecology*, *20*(6), 703–717. <https://doi.org/10.1007/s10980-005-0063-7>
- Graham, M. H. (2003). Confronting multicollinearity in ecological multiple regression. *Ecology*, *84*(11), 2809–2815. <https://doi.org/10.1890/02-3114>
- Guisan, A., Thuiller, W., & Zimmermann, N. E. (2017). *Habitat suitability and distribution models: With applications in R*. Cambridge, UK: Cambridge University Press. <https://doi.org/10.1017/9781139028271>
- Guo, F., Lenoir, J., & Bonebrake, T. C. (2018). Land-use change interacts with climate to determine elevational species redistribution. *Nature Communications*, *9*(1), 1315. <https://doi.org/10.1038/s41467-018-03786-9>
- Guo, X., Feng, J., Shi, Z., Zhou, X., Yuan, M., Tao, X., ... Zhou, J. (2018). Climate warming leads to divergent succession of grassland microbial communities. *Nature Climate Change*, *8*(9), 813–818. <https://doi.org/10.1038/s41558-018-0254-2>
- Hagedorn, F., Gavazov, K., & Alexander, J. M. (2019). Above- and below-ground linkages shape responses of mountain vegetation to climate change. *Science*, *365*(6458), 1119–1123. <https://doi.org/10.1126/science.aax4737>
- Hannemann, H., Willis, K. J., & Macias-Fauria, M. (2016). The devil is in the detail: Unstable response functions in species distribution models challenge bulk ensemble modelling. *Global Ecology and Biogeography*, *25*(1), 26–35. <https://doi.org/10.1111/geb.12381>
- Hao, T. X., Elith, J., Guillerá-Arroita, G., & Lahoz-Monfort, J. J. (2019). A review of evidence about use and performance of species distribution modelling ensembles like BIOMOD. *Diversity and Distributions*, *25*(5), 839–852. <https://doi.org/10.1111/ddi.12892>
- Harrison, S. P., Prentice, I. C., Barboni, D., Kohfeld, K. E., Ni, J., & Sutra, J.-P. (2010). Ecophysiological and bioclimatic foundations for a global plant functional classification. *Journal of Vegetation Science*, *21*(2), 300–317. <https://doi.org/10.1111/j.1654-1103.2009.01144.x>
- Hirzel, A., & Guisan, A. (2002). Which is the optimal sampling strategy for habitat suitability modelling. *Ecological Modelling*, *157*(2), 331–341. [https://doi.org/10.1016/S0304-3800\(02\)00203-X](https://doi.org/10.1016/S0304-3800(02)00203-X)
- Illan, J. G., Gutierrez, D., & Wilson, R. J. (2010). The contributions of topoclimate and land cover to species distributions and abundance: Fine-resolution tests for a mountain butterfly fauna. *Global Ecology and Biogeography*, *19*(2), 159–173. <https://doi.org/10.1111/j.1466-8238.2009.00507.x>
- IPBES. (2019). *Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science – Policy Platform on Biodiversity and Ecosystem Services*. In E. S. Brondizio, J. Settele, S. Díaz, & H. T. Ngo (Eds.). Bonn, Germany: IPBES Secretariat.
- IPCC. (2014). *Climate change 2014: Synthesis report. Contribution of working groups I, II and III to the fifth assessment report of the Intergovernmental Panel on Climate Change*. In Core Writing Team, R. K. Pachauri, & L. A. Meyer (Eds.). Geneva, Switzerland: Intergovernmental Panel on Climate Change. Retrieved from [https://www.ipcc.ch/site/assets/uploads/2018/05/SYR\\_AR5\\_FINAL\\_full\\_wcover.pdf](https://www.ipcc.ch/site/assets/uploads/2018/05/SYR_AR5_FINAL_full_wcover.pdf)
- Jiménez-Valverde, A., & Lobo, J. M. (2007). Threshold criteria for conversion of probability of species presence to either-or presence-absence. *Acta Oecologica*, *31*(3), 361–369. <https://doi.org/10.1016/j.actao.2007.02.001>
- Karger, D. N., Conrad, O., Böhrner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., ... Kessler, M. (2017). Climatologies at high resolution for the earth's land surface areas. *Scientific Data*, *4*, 170122. <https://doi.org/10.1038/sdata.2017.122>
- Kharouba, H. M., McCune, J. L., Thuiller, W., & Huntley, B. (2013). Do ecological differences between taxonomic groups influence the relationship between species' distributions and climate? A global meta-analysis using species distribution models. *Ecography*, *36*(6), 657–664. <https://doi.org/10.1111/j.1600-0587.2012.07683.x>
- Lawrence, D. J., Stewart-Koster, B., Olden, J. D., Ruesch, A. S., Torgersen, C. E., Lawler, J. J., ... Crown, J. K. (2014). The interactive effects of climate change, riparian management, and a nonnative predator on stream-rearing salmon. *Ecological Applications*, *24*(4), 895–912. <https://doi.org/10.1890/13-0753.1>
- Lawson, C. R., Hodgson, J. A., Wilson, R. J., & Richards, S. A. (2014). Prevalence, thresholds and the performance of presence-absence models. *Methods in Ecology and Evolution*, *5*(1), 54–64. <https://doi.org/10.1111/2041-210X.12123>
- Lembrechts, J. J., Aalto, J., Ashcroft, M. B., De Frenne, P., Kopecký, M., Lenoir, J., ... Nijs, I. (2020). SoilTemp: A global database of near-surface temperature. *Global Change Biology*. <https://doi.org/10.1111/gcb.15123>
- Lembrechts, J. J., Lenoir, J., Roth, N., Hattab, T., Milbau, A., Haider, S., ... Nijs, I. (2019). Comparing temperature data sources for use in

- species distribution models: From in-situ logging to remote sensing. *Global Ecology and Biogeography*, 28(11), 1578–1596. <https://doi.org/10.1111/geb.12974>
- Lembrechts, J. J., Nijs, I., & Lenoir, J. (2019). Incorporating microclimate into species distribution models. *Ecography*, 42(7), 1267–1279. <https://doi.org/10.1111/ecog.03947>
- Lenoir, J., Bertrand, R., Comte, L., Bourgeaud, L., Hattab, T., Murielle, J., & Grenouillet, G. (2020). Species better track climate warming in the oceans than on land. *Nature Ecology & Evolution*, 4(8), 1044–1059. <https://doi.org/10.1038/s41559-020-1198-2>
- Lenoir, J., Gégout, J.-C., Marquet, P. A., de Ruffray, P., & Brisse, H. (2008). A significant upward shift in plant species optimum elevation during the 20th century. *Science*, 320(5884), 1768–1771. <https://doi.org/10.1126/science.1156831>
- Lenoir, J., Graae, B. J., Aarrestad, P. A., Alsos, I. G., Armbruster, W. S., Austrheim, G., ... Svenning, J.-C. (2013). Local temperatures inferred from plant communities suggest strong spatial buffering of climate warming across Northern Europe. *Global Change Biology*, 19(5), 1470–1481. <https://doi.org/10.1111/gcb.12129>
- Lenoir, J., Hattab, T., & Pierre, G. (2017). Climatic microrefugia under anthropogenic climate change: Implications for species redistribution. *Ecography*, 40(2), 253–266. <https://doi.org/10.1111/ecog.02788>
- Lenoir, J., & Svenning, J.-C. (2015). Climate-related range shifts – A global multidimensional synthesis and new research directions. *Ecography*, 38(1), 15–28. <https://doi.org/10.1111/ecog.00967>
- Li, S., Shakoor, A., Wubet, T., Zhang, N. L., Liang, Y., & Ma, K. P. (2018). Fine-scale variations of fungal community in a heterogeneous grassland in Inner Mongolia: Effects of the plant community and edaphic parameters. *Soil Biology & Biochemistry*, 122, 104–110. <https://doi.org/10.1016/j.soilbio.2018.04.007>
- Liu, C., Berry, P. M., Dawson, T. P., & Pearson, R. G. (2005). Selecting thresholds of occurrence in the prediction of species distributions. *Ecography*, 28(3), 385–393. <https://doi.org/10.1111/j.0906-7590.2005.03957.x>
- Lobo, J. M., Jiménez-Valverde, A., & Real, R. (2008). AUC: A misleading measure of the performance of predictive distribution models. *Global Ecology and Biogeography*, 17(2), 145–151. <https://doi.org/10.1111/j.1466-8238.2007.00358.x>
- Lomba, A., Pellissier, L., Randin, C., Vicente, J., Moreira, F., Honrado, J., & Guisan, A. (2010). Overcoming the rare species modelling paradox: A novel hierarchical framework applied to an Iberian endemic plant. *Biological Conservation*, 143(11), 2647–2657. <https://doi.org/10.1016/j.biocon.2010.07.007>
- Malard, L. A., Anwar, M. Z., Jacobsen, C. S., & Pearce, D. A. (2019). Biogeographical patterns in soil bacterial communities across the Arctic region. *FEMS Microbiology Ecology*, 95(9). <https://doi.org/10.1093/femsec/fiz128>
- McGraw, K. O., & Wong, S. P. (1992). A common language effect size statistic. *Psychological Bulletin*, 111(2), 361–365. <https://doi.org/10.1037/0033-2909.111.2.361>
- Menéndez, R., González-Megías, A., Jay-Robert, P., & Marquéz-Ferrando, R. (2014). Climate change and elevational range shifts: Evidence from dung beetles in two European mountain ranges. *Global Ecology and Biogeography*, 23(6), 646–657. <https://doi.org/10.1111/geb.12142>
- Miller, D. A. W., Grant, E. H. C., Muths, E., Amburgey, S. M., Adams, M. J., Joseph, M. B., ... Sigafus, B. H. (2018). Quantifying climate sensitivity and climate-driven change in North American amphibian communities. *Nature Communications*, 9(1), 3926. <https://doi.org/10.1038/s41467-018-06157-6>
- Mod, H. K., & Luoto, M. (2016). Arctic shrubification mediates the impacts of warming climate on changes to tundra vegetation. *Environmental Research Letters*, 11(12), 124028. <https://doi.org/10.1088/1748-9326/11/12/124028>
- Mod, H. K., Scherrer, D., Luoto, M., & Guisan, A. (2016). What we use is not what we know: Environmental predictors in plant distribution models. *Journal of Vegetation Science*, 27(6), 1308–1322. <https://doi.org/10.1111/jvs.12444>
- Moeslund, J. E., Arge, L., Bøcher, P. K., Dalgaard, T., Ejrnæs, R., Odgaard, M. V., & Svenning, J.-C. (2013). Topographically controlled soil moisture drives plant diversity patterns within grasslands. *Biodiversity and Conservation*, 22(10), 2151–2166. <https://doi.org/10.1007/s10531-013-0442-3>
- Montgomery, K. (2006). Variation in temperature with altitude and latitude. *Journal of Geography*, 105(3), 133–135. <https://doi.org/10.1080/00221340608978675>
- Moritz, C., & Agudo, R. (2013). The future of species under climate change: Resilience or decline? *Science*, 341(6145), 504–508. <https://doi.org/10.1126/science.1237190>
- Nenzen, H. K., & Araujo, M. B. (2011). Choice of threshold alters projections of species range shifts under climate change. *Ecological Modelling*, 222(18), 3346–3354. <https://doi.org/10.1016/j.ecolmodel.2011.07.011>
- Niittynen, P., Heikkinen, R. K., & Luoto, M. (2018). Snow cover is a neglected driver of Arctic biodiversity loss. *Nature Climate Change*, 8(11), 997–1001. <https://doi.org/10.1038/s41558-018-0311-x>
- Oliver, T. H., & Morecroft, M. D. (2014). Interactions between climate change and land use change on biodiversity: Attribution problems, risks, and opportunities. *WIREs Climate Change*, 5(3), 317–335. <https://doi.org/10.1002/wcc.271>
- Pacifici, M., Foden, W. B., Visconti, P., Watson, J. E. M., Butchart, S. H. M., Kovacs, K. M., ... Rondinini, C. (2015). Assessing species vulnerability to climate change. *Nature Climate Change*, 5, 215–224. <https://doi.org/10.1038/nclimate2448>
- Pagni, M., Niculita-Hirzel, H., Pellissier, L., Dubuis, A., Xenarios, I., Guisan, A., ... Guex, N. (2013). Density-based hierarchical clustering of pyrosequences on a large scale – The case of fungal ITS1. *Bioinformatics*, 29(10), 1268–1274. <https://doi.org/10.1093/bioinformatics/btt149>
- Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, 421(6918), 37–42. <https://doi.org/10.1038/nature01286>
- Peay, K. G., Kennedy, P. G., & Talbot, J. M. (2016). Dimensions of biodiversity in the Earth mycobiome. *Nature Reviews Microbiology*, 14(7), 434–447. <https://doi.org/10.1038/nrmicro.2016.59>
- Pellissier, L., Pradervand, J.-N., Pottier, J., Dubuis, A., Maiorano, L., & Guisan, A. (2012). Climate-based empirical models show biased predictions of butterfly communities along environmental gradients. *Ecography*, 35(8), 684–692. <https://doi.org/10.1111/j.1600-0587.2011.07047.x>
- Pellissier, L., Pradervand, J.-N., Williams, P. H., Litsios, G., Cherix, D., Guisan, A., & Baselga, A. (2013). Phylogenetic relatedness and proboscis length contribute to structuring bumblebee communities in the extremes of abiotic and biotic gradients. *Global Ecology and Biogeography*, 22(5), 577–585. <https://doi.org/10.1111/geb.12026>
- Petitpierre, B., McDougall, K., Seipel, T., Broennimann, O., Guisan, A., & Kueffer, C. (2016). Will climate change increase the risk of plant invasions into mountains? *Ecological Applications*, 26(2), 530–544. <https://doi.org/10.1890/14-1871>
- Pinol, J., Mir, G., Gomez-Polo, P., & Agusti, N. (2015). Universal and blocking primer mismatches limit the use of high-throughput DNA sequencing for the quantitative metabarcoding of arthropods. *Molecular Ecology Resources*, 15(4), 819–830. <https://doi.org/10.1111/1755-0998.12355>
- Pinto-Figueroa, E. A., Seddon, E., Yashiro, E., Buri, A., Niculita-Hirzel, H., van der Meer, J. R., & Guisan, A. (2019). Archaeorhizomycetes spatial distribution in soils along wide elevational and environmental gradients reveal co-abundance patterns with other fungal saprobes and potential weathering capacities. *Frontiers in Microbiology*, 10(656), 656. <https://doi.org/10.3389/fmicb.2019.00656>
- Pittet, M. (2017). *Impact of global warming on the distribution and dispersal of reptiles in the Western Swiss Alps*. Master of Science, Université de Lausanne, Lausanne.

- Porter, T. M., & Hajibabaei, M. (2018). Scaling up: A guide to high-throughput genomic approaches for biodiversity analysis. *Molecular Ecology*, 27(2), 313–338. <https://doi.org/10.1111/mec.14478>
- Pradervand, J.-N., Dubuis, A., Reymond, A., Sonnay, V., Gelin, A., & Guisan, A. (2013). Quel facteur influencent la richesse en orthoptères des Préalpes vaudoises? *Bulletin de la Société Vaudoise des Sciences Naturelles*, 93(4), 155–173.
- Randin, C. F., Dirnböck, T., Dullinger, S., Zimmermann, N. E., Zappa, M., & Guisan, A. (2006). Are niche-based species distribution models transferable in space? *Journal of Biogeography*, 33(10), 1689–1703. <https://doi.org/10.1111/j.1365-2699.2006.01466.x>
- Riesenfeld, C. S., Schloss, P. D., & Handelsman, J. (2004). Metagenomics: Genomic analysis of microbial communities. *Annual Review of Genetics*, 38(1), 525–552. <https://doi.org/10.1146/annurev.genet.38.072902.091216>
- Roberts, D. R., Nielsen, S. E., & Stenhouse, G. B. (2014). Idiosyncratic responses of grizzly bear habitat to climate change based on projected food resource changes. *Ecological Applications*, 24(5), 1144–1154. <https://doi.org/10.1890/13-0829.1>
- Rodríguez-Castañeda, G., Hof, A. R., Jansson, R., & Harding, L. E. (2012). Predicting the fate of biodiversity using species' distribution models: Enhancing model comparability and repeatability. *PLoS One*, 7(9), e44402. <https://doi.org/10.1371/journal.pone.0044402>
- Rogelj, J., Meinshausen, M., & Knutti, R. (2012). Global warming under old and new scenarios using IPCC climate sensitivity range estimates. *Nature Climate Change*, 2(4), 248–253. <https://doi.org/10.1038/nclimate1385>
- Scherrer, D., & Guisan, A. (2019). Ecological indicator values reveal missing predictors of species distributions. *Scientific Reports*, 9(1), 3061. <https://doi.org/10.1038/s41598-019-39133-1>
- Scherrer, D., & Körner, C. (2011). Topographically controlled thermal-habitat differentiation buffers alpine plant diversity against climate warming. *Journal of Biogeography*, 38(2), 406–416. <https://doi.org/10.1111/j.1365-2699.2010.02407.x>
- Scherrer, D., Mod, H. K., & Guisan, A. (2020). How to evaluate community predictions without thresholding? *Methods in Ecology and Evolution*, 11(1), 51–63. <https://doi.org/10.1111/2041-210x.13312>
- Scherrer, D., Schmid, S., & Körner, C. (2011). Elevational species shifts in a warmer climate are overestimated when based on weather station data. *International Journal of Biometeorology*, 55(4), 645–654. <https://doi.org/10.1007/s00484-010-0364-7>
- Schleuning, M., Fründ, J., Schweiger, O., Welk, E., Albrecht, J., Albrecht, M., ... Hof, C. (2016). Ecological networks are more sensitive to plant than to animal extinction under climate change. *Nature Communications*, 7, 13965. <https://doi.org/10.1038/ncomms13965>
- Schmidt, B. R., & Zumbach, S. (2019). Amphibian conservation in Switzerland. In H. Heatwole & J. W. Wilkinson (Eds.), *Status of conservation and decline of amphibians: Eastern hemisphere* (Vol. 5, pp. 46–51). Exeter: Pelagic Publishing.
- Sears, M. W., Raskin, E., & Angilletta Jr., M. J. (2011). The world is not flat: Defining relevant thermal landscapes in the context of climate change. *Integrative and Comparative Biology*, 51(5), 666–675. <https://doi.org/10.1093/icb/acr111>
- Segurado, P., Santos, J. M., Pont, D., Melcher, A. H., Jalon, D. G., Hughes, R. M., & Ferreira, M. T. (2011). Estimating species tolerance to human perturbation: Expert judgment versus empirical approaches. *Ecological Indicators*, 11(6), 1623–1635. <https://doi.org/10.1016/j.ecolind.2011.04.006>
- Seibold, S., Cadotte, M. W., Maclvor, J. S., Thorn, S., & Muller, J. (2018). The necessity of multitrophic approaches in community ecology. *Trends in Ecology & Evolution*, 33(10), 754–764. <https://doi.org/10.1016/j.tree.2018.07.001>
- Seoane, J., Bustamante, J., & Diaz-Delgado, R. (2004). Competing roles for landscape, vegetation, topography and climate in predictive models of bird distribution. *Ecological Modelling*, 171(3), 209–222. <https://doi.org/10.1016/j.ecolmodel.2003.08.006>
- Seppey, C. V. W., Broennimann, O., Buri, A., Yashiro, E., Pinto-Figueroa, E., Singer, D., ... Lara, E. (2020). Soil protist diversity in the Swiss western Alps is better predicted by topo-climatic than by edaphic variables. *Journal of Biogeography*, 47(4), 866–878. <https://doi.org/10.1111/jbi.13755>
- Serna-Chavez, H. M., Fierer, N., & van Bodegom, P. M. (2013). Global drivers and patterns of microbial abundance in soil. *Global Ecology and Biogeography*, 22(10), 1162–1172. <https://doi.org/10.1111/geb.12070>
- Somodi, I., Lepesi, N., & Botta-Dukat, Z. (2017). Prevalence dependence in model goodness measures with special emphasis on true skill statistics. *Ecology and Evolution*, 7(3), 863–872. <https://doi.org/10.1002/ece3.2654>
- Soudzilovskaia, N. A., Elumeeva, T. G., Onipchenko, V. G., Shidakov, I. I., Salpagarova, F. S., Khubiev, A. B., ... Cornelissen, J. H. C. (2013). Functional traits predict relationship between plant abundance dynamic and long-term climate warming. *Proceedings of the National Academy of Sciences of the United States of America*, 110(45), 18180–18184. <https://doi.org/10.1073/pnas.1310700110>
- Staniczenko, P. P. A., Sivasubramaniam, P., Suttle, K. B., & Pearson, R. G. (2017). Linking macroecology and community ecology: Refining predictions of species distributions using biotic interaction networks. *Ecology Letters*, 20(6), 693–707. <https://doi.org/10.1111/ele.12770>
- Steidinger, B. S., Crowther, T. W., Liang, J., Van Nuland, M. E., Werner, G. D. A., Reich, P. B., ... Peay, K. G. (2019). Climatic controls of decomposition drive the global biogeography of forest-tree symbioses. *Nature*, 569(7756), 404–408. <https://doi.org/10.1038/s41586-019-1128-0>
- Steinbauer, M. J., Grytnes, J.-A., Jurasinski, G., Kulonen, A., Lenoir, J., Pauli, H., ... Wipf, S. (2018). Accelerated increase in plant species richness on mountain summits is linked to warming. *Nature*, 556(7700), 231–234. <https://doi.org/10.1038/s41586-018-0005-6>
- Suggitt, A. J., Gillingham, P. K., Hill, J. K., Huntley, B., Kunin, W. E., Roy, D. B., & Thomas, C. D. (2011). Habitat microclimates drive fine-scale variation in extreme temperatures. *Oikos*, 120(1), 1–8. <https://doi.org/10.1111/j.1600-0706.2010.18270.x>
- Tasser, E., & Tappeiner, U. (2002). Impact of land use changes on mountain vegetation. *Applied Vegetation Science*, 5(2), 173–184. <https://doi.org/10.1111/j.1654-109X.2002.tb00547.x>
- Tedersoo, L., Bahram, M., Pölme, S., Kõljalg, U., Yorou, N. S., Wijesundera, R., ... Abarenkov, K. (2014). Global diversity and geography of soil fungi. *Science*, 346(6213), 1256688. <https://doi.org/10.1126/science.1256688>
- Thuiller, W., Georges, D., Engler, R., & Breiner, F. (2016). biomod2: Ensemble platform for species distribution modeling R-package (version 3.3-7). Retrieved from <https://CRAN.R-project.org/package=biomod2>
- Thuiller, W., Lafourcade, B., Engler, R., & Araújo, M. B. (2009). BIOMOD – A platform for ensemble forecasting of species distributions. *Ecography*, 32(3), 369–373. <https://doi.org/10.1111/j.1600-0587.2008.05742.x>
- Troudet, J., Grandcolas, P., Blin, A., Vignes-Lebbe, R., & Legendre, F. (2017). Taxonomic bias in biodiversity data and societal preferences. *Scientific Reports*, 7(1), 9132. <https://doi.org/10.1038/s41598-017-09084-6>
- Vaughan, I. P., & Ormerod, S. J. (2005). The continuing challenges of testing species distribution models. *Journal of Applied Ecology*, 42(4), 720–730. <https://doi.org/10.1111/j.1365-2664.2005.01052.x>
- Walther, G.-R. (2010). Community and ecosystem responses to recent climate change. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1549), 2019–2024. <https://doi.org/10.1098/rstb.2010.0021>

- Wang, L., Qiu, X., Wang, P., Wang, X., & Liu, A. (2014). Influence of complex topography on global solar radiation in the Yangtze River Basin. *Journal of Geographical Sciences*, 24(6), 980–992. <https://doi.org/10.1007/s11442-014-1132-0>
- Wardle, D. A., Bardgett, R. D., Klironomos, J. N., Setälä, H., van der Putten, W. H., & Wall, D. H. (2004). Ecological linkages between aboveground and belowground biota. *Science*, 304(5677), 1629–1633. <https://doi.org/10.1126/science.1094875>
- Warren, D. L., Glor, R. E., & Turelli, M. (2008). Environmental niche equivalency versus conservatism: Quantitative approaches to niche evolution. *Evolution*, 62(11), 2868–2883. <https://doi.org/10.1111/j.1558-5646.2008.00482.x>
- Wilkinson, D. M. (1998). Fragments of an entangled bank: Do ecologists study most of ecology? *Oikos*, 82(2), 393–394. <https://doi.org/10.2307/3546981>
- Woodward, F. I., & Williams, B. G. (1987). Climate and plant distribution at global and local scales. *Vegetatio*, 69(1), 189–197. <https://doi.org/10.1007/bf00038700>
- Yashiro, E., Pinto-Figueroa, E., Buri, A., Spangenberg, J. E., Adatte, T., Niculita-Hirzel, H., ... van der Meer, J. R. (2016). Local environmental factors drive divergent grassland soil bacterial communities in the Western Swiss Alps. *Applied and Environmental Microbiology*, 82(21), 6303–6316. <https://doi.org/10.1128/AEM.01170-16>
- Yashiro, E., Pinto-Figueroa, E., Buri, A., Spangenberg, J. E., Adatte, T., Niculita-Hirzel, H., ... van der Meer, J. R. (2018). Meta-scale mountain grassland observatories uncover commonalities as well as specific interactions among plant and non-rhizosphere soil bacterial communities. *Scientific Reports*, 8(1), 5758. <https://doi.org/10.1038/s41598-018-24253-x>
- Yoccoz, N. G. (2012). The future of environmental DNA in ecology. *Molecular Ecology*, 21(8), 2031–2038. <https://doi.org/10.1111/j.1365-294X.2012.05505.x>
- Zubler, E. M., Fischer, A. M., Liniger, M. A., Croci-Maspoli, M., Scherrer, S. C., & Appenzeller, C. (2014). Localized climate change scenarios of mean temperature and precipitation over Switzerland. *Climatic Change*, 125(2), 237–252. <https://doi.org/10.1007/s10584-014-1144-x>

#### SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Mod HK, Scherrer D, Di Cola V, et al. Greater topoclimatic control of above- versus below-ground communities. *Glob Change Biol*. 2020;26:6715–6728. <https://doi.org/10.1111/gcb.15330>