



Det helsevitenskapelige fakultet
Institutt for samfunnsmedisin

ILSA 2017 in Tromsø

*Proceedings from the 42nd annual conference of the
International Lung Sound Association*

—
edited by Hasse Melbye

ISM skriftserie nr. 182



*ISM skriftserie
blir utgitt av Institutt for samfunnsmedisin,
UiT Norges arktiske universitet.*

*Forfatterne er selv ansvarlige for sine funn og
konklusjoner. Innholdet er derfor ikke uttrykk
for ISM's syn.*

*The opinions expressed in this publication are those
of the authors and do not necessarily reflect the
official policy of the institutions supporting this research.*

ISBN: 978 - 82 - 8378 -003 - 1
2017

Table of contents

(presenting authors in brackets)

History of ILSA conferences	7
The 42 nd ILSA conference	9
Abstracts:	
Lung sound detection and monitoring	
Prevalence and associations of wheezes and crackles in a general adult population - the Tromsø 7 study. <i>(Juan Carlos Aviles Solis)</i>	11
Reliability and validity of computerized respiratory sounds in COPD <i>(Ana Oliveira)</i>	12
Monitoring of nocturnal respiratory symptoms in COPD patients by using LEOSound monitor <i>(Patrick Fischer)</i>	13
Lung sounds intensity: is there a difference between spontaneous and target airflow? <i>(Cristina Jácome)</i>	14
The influence of adventitious sounds and artifacts on the frequency and intensity of normal respiratory sounds <i>(Cristina Jácome)</i>	
Machine learning	
From another research field: Using machine learning to provide automatic image annotation for wildlife camera traps in the Arctic <i>(John Markus Bjørndalen)</i>	17
A New Computerized Approach for Lung Sounds Analysis Using Sound Signature <i>(Doran Adler)</i>	21
Use Wide Range Sound Signature in Computerized Classification of Lung Diseases <i>(Doran Adler)</i>	23
Detection of Wheezes and Breathing Phases Using Deep Convolutional Neural Networks <i>(Johan Ravn)</i>	24
Machine Learning Based Crackle Detection <i>(Morten Grønnesby)</i>	27
Comparison of Various Machine Learning Algorithms on Adventitious Pulmonary Sound Classification <i>(Yasemin Kahya)</i>	28
Recording in the clinic	
Cardiac Response to Respiration in Deep Inspiration followed by Breath Holding <i>(Sadamu Ishikawa)</i>	29
Auscultatory phenomena in patients with partially controlled bronchial asthma with intrapulmonary auscultation <i>(Yuriy Petrov)</i>	30
Lung sounds of patients with interstitial pneumonia in left and right lateral decubitus position – Effect of gravity on lung sounds <i>(Michico Tsuchiya)</i>	31
Comparing lung sounds intensity between sitting, supine and prone position in patients with interstitial pneumonia <i>(Michico Tsuchiya)</i>	33
Lung sound changes associated with pneumothorax <i>(Hansen Mansy)</i>	35
Crackles and Rhonchi with Postural Dependence in a 9-year-old Girl with Severe Mycoplasmal Pneumonia <i>(Masato Takase)</i>	36
Rumbling rhonchi in bronchial asthma: Their relation to airway inflammation <i>(Yukio Nagasaka)</i>	37

Rumbling rhonchi and bronchial inflammation in patients with bronchial asthma (<i>Yukio Nagasaka</i>)	38
Lung sounds of a patient with bronchiolitis obliterans: comparison with pulmonary emphysema (<i>Hiroshi Nakano</i>)	40
Lung sounds in clinical diagnosis	
Sub-classification of wheezes and crackles is not helpful when lung auscultation is used for detecting decreased lung function in adults (<i>Hasse Melbye</i>)	42
Can normal respiratory sounds differentiate COPD grades? (<i>Alda Marques</i>)	44
Is Auscultation Still Useful or is it Just Nostalgia? (<i>Steve Kraman</i>)	45
Lung auscultation - psychology and learning	
Psychoacoustics of auscultation (<i>Hans Pasterkamp</i>)	48
Teaching and Learning pulmonary auscultation – the Computerized Lung Auscultation – Sound Software (CLASS) (<i>Alda Marques</i>)	50
Improved classification of lung sounds by medical students when spectrograms are displayed (<i>Juan Carlos Aviles Solis</i>)	52
Making an illustrated lung auscultation guide book for nurses: are these illustrations appropriate? (<i>Yukio Nagasaka</i>)	53
Live Demonstration of an Advanced Pulmonary Auscultation Simulator (<i>Thomas Talbot</i>)	55
Various topics	
Pulmonary Rehabilitation effects on computerized respiratory sounds of patients with AECOPD (<i>Alda Marques</i>)	57
New features of 10-19 kHz sound propagation through human lungs (<i>Alexander Dyachenko</i>)	59
Differences in waveforms and power spectra of chest wall percussion sounds obtained by various sensors (<i>Alexander Dyachenko</i>)	61
Using cough epochs to describe nocturnal cough in COPD patients (<i>Patrick Fischer</i>)	62
Investigation of pre- and postoperative lung sounds of patients who underwent general thoracic surgery (<i>Takashi Suzuki</i>)	63
National survey on the perception of respiratory sounds among Greek doctors (preliminary report) (<i>Kostas Priftis</i>)	64

History of ILSA conferences

In October 1976, the First International Conference on Lung Sounds was held in Boston, MA. The objectives of this conference were defined as follows:

"Studies of lung sounds have been reported with increasing frequency in recent years. This conference is convened to provide an opportunity for exchange of ideas and experience among those who have an active interest in the subject. Clinicians, physiologists, engineers, and perceptual psychologists can each contribute towards a better understanding of what lung sounds mean. They will have a better chance of doing so after talking together".

"We hope that comparisons of methods of recording, analyzing, and describing lung sounds will reduce ambiguity. We hope that discussions about work in progress may prevent unnecessary duplication of effort. We hope that investigators will save time and avoid some mistakes by learning what others have done".

ILSA conferences have been arranged every year since then, driven by research interest among medical doctors, engineers and researchers from other relevant fields, together with the fast development of technology for recording and analyzing lung sounds.

No.	Date	Place	Local Organizer(s)
1	October, 1976	Boston, MA	Raymond L.H. Murphy, Jr.
2	September, 1977	Cincinnati, OH	Robert Loudon
3	September, 1978	New Orleans, LA	William Waring
4	September, 1979	Chicago, IL	David Cugell
5	September, 1980	London, England	Leslie Capel & Paul Forgacs
6	October, 1981	Boston, MA	Raymond L.H. Murphy, Jr.
7	October, 1982	Martinez, CA	Peter Krumpal
8	September, 1983	Baltimore, MD	Wilmot Ball
9	September, 1984	Cincinnati, OH	Robert Loudon
10	September, 1985	Tokyo, Japan	Riichiro Mikami
11	September, 1986	Lexington, KY	Steve S. Kraman
12	September, 1987	Paris, France	Gerard Charbonneau
13	September, 1988	Chicago, IL	David Cugell
14	September, 1989	Winnipeg, Canada	Hans Pasterkamp
15	October, 1990	New Orleans, LA	David Rice
16	September, 1991	Veruno, Italy	Filiberto Dalmaso

17	August, 1992	Helsinki, Finland	Anssi Sovijarvi
18	August, 1993	Alberta, Canada	Raphael Beck
19	September, 1994	Haifa, Israel	Noam Gavriely
20	October, 1995	Long Beach, CA	Christopher Druzgalski
21	September, 1996	Chester, England	John Earis
22	October, 1997	Tokyo, Japan	Masashi Mori
23	October, 1998	Boston, MA	Sadamu Ishikawa
24	October, 1999	Marburg, Germany	Peter von Wichert
25	September, 2000	Chicago, IL	David Cugell
26	September, 2001	Berlin, Germany	Hans Pasterkamp
27	September, 2002	Helsinki, Finland	Anssi Sovijarvi
28	September, 2003	Cancun, Mexico	Sonia Charleston, Ramon Gonzales Camerena & Tomas Aljama Corrales
29	September, 2004	Glasgow, Scotland	Ken Anderson & John Earis
30	September, 2005	Boston, MA	Raymond L.H. Murphy, Jr.
31	September, 2006	Halkidiki, Greece	Leontios Hadjileontiadis
32	November, 2007	Tokyo, Japan	Shoji Kudoh
33	October, 2008	Boston, MA	Sadamu Ishikawa & Raymond L.H. Murphy, Jr.
34	September, 2009	Haifa, Israel	Noam Gavriely
35	October, 2010	Toledo, Ohio	Dan E. Olson
36	September, 2011	Manchester, England	Ashley Woodcock
37	October, 2012	Mayo Clinic, MN	Rasanen Jukka
38	November, 2013	Kyoto, Japan	Yukio Nagasaka
39	October, 2014	Boston, MA	Sadamu Ishikawa
40	September, 2015	Saint-Petersburg, Russia	Alexander Dyachenko
41	October, 2016	Tokyo, Japan	Masato Takase & Shoji Kudoh
42	September, 2017	Tromsø, Norway	Hasse Melbye



The 42nd ILSA conference in Tromsø

The usefulness of lung auscultation is changing. It depends on how well practitioners understand the generation of sounds. It also depends on their knowledge on how lung sounds are associated with lung and heart diseases, as well as with other factors such as ageing and smoking habits. In clinical practice, practitioners need to give sufficient attention to lung auscultation, and they should use the same terminology, or at least understand each other's use of terms. Technological innovations lead to an extended use of lung auscultation. Continuous monitoring of lung sounds is now possible, and computers can extract more information from the complex lung sounds than human hearing is capable of. Learning how to carry out lung auscultation and to interpret the sounds are essential skills in the education of doctors and other health professionals. Thus, new computer based learning tools for the study of recorded sounds will be helpful.

In this conference there will be focus on all these determinants for efficient lung auscultation. In addition to free oral presentations, we have three symposia: on computerized analysis based on machine learning, on diagnostics, and on learning lung sounds, including the psychology of hearing. The symposia include extended presentations from invited speakers.

The 42nd conference is the first in history arranged by a research unit for general practice. Primary care doctors are probably the group of health professionals that put the greatest emphasis on lung auscultation in their clinical work. Many patients with chest symptoms consult without a known diagnosis, and several studies have shown that general practitioners pay attention to crackles and wheezes when making decisions, for instance when antibiotics are prescribed to coughing patients. In hospital, the diagnosis of lung diseases is more strongly influenced by technologies such as radiography and blood gas analysis. Since lung auscultation holds a strong position in the work of primary care doctors, I think it is just timely, that the 42nd ILSA conference is hosted by General Practice Research Unit in Tromsø. I hope all participants will find presentations of importance, and that the stay in Tromsø will be enjoyable.

Hasse Melbye, conference president (hasse.melbye@uit.no)

Professor at General Practice Research Unit, Department of Community Medicine, Faculty of Health Sciences, UIT The Arctic University of Norway

Prevalence and associations of crackles and wheezes in a general adult population - the Tromsø 7 study


Aviles-Solis JC¹ (juan.c.solis@uit.no), Jácome C.^{1,2}, Pasterkamp H.³, and Melbye H.¹

¹ Department of Community Medicine, UIT the Arctic University of Norway. ² Lab3R - Respiratory Research and Rehabilitation Laboratory, School of Health Sciences, University of Aveiro. ³ Dept. of Pediatrics and Child Health, University of Manitoba, Canada.

Background: The stethoscope is a medical tool that is broadly available, cheap, and easy to use. However, there is a lack of epidemiological data about abnormal lung sounds in relation to chronic heart and lung diseases.

Aims: To explore the prevalence of crackles and wheezes in a general population and detect explanatory factors

Methods: We recorded respiratory sounds from a random sample of 4,033 participants from the Tromsø 7 study (n=21,083), using a microphone set in a stethoscope tube. We obtained recordings from six locations on the chest (figure). The participants were 40 to 84 years old. In addition, we obtained information on symptoms, self-reported disease and clinical tests. Four observers classified the recordings independently, and several rounds of classification were carried out to optimize reliability. Chi-square tests were used to explore associations.



Results: We found adventitious sounds in 28% (n=1131) of the individuals, of which 13% (n=532) presented with crackles and 18% (n=729) with wheezes. In about 71.6% (crackles) and 73.25% (wheezes) of the individuals, the positive finding was present in only one of the six locations. The prevalence of adventitious sounds was evenly distributed across the six chest locations except for inspiratory crackles that were more frequent in the bases of the lungs. Daily cough ($p < 0.001$) and Modified Medical Research Council (mMRC) score > 2 ($p < 0.001$) were associated with inspiratory crackles. Present (but not past) smoking was associated with both crackles ($p < 0.001$) and wheezes ($p < 0.01$). Self-reported heart attack ($p < 0.01$), COPD ($p < 0.001$) and rheumatoid arthritis ($p < 0.001$) were also associated with crackles.

Conclusions: The presence of adventitious lung sounds was common in our sample. We found an association of positive findings with respiratory symptoms, smoking status and self-reported disease. Further analyses will explore the relationship between crackles and wheezes with defined diagnostic categories for heart and lung diseases.

Reliability and validity of computerised respiratory sounds in COPD

Ana Oliveira¹⁻³ (alao@ua.pt) Susan Lage⁴, Joana Neves⁵ Alda Marques^{2,3}

¹Faculty of Sports, University of Porto, Portugal, ²Lab 3R – Respiratory Research and Rehabilitation Laboratory, School of Health Sciences, University of Aveiro, Portugal

³Institute for Research in Biomedicine (iBiMED), University of Aveiro, Portugal

⁴Rehabilitation Science Program, Federal University of Minas Gerais, Brasil

⁵Centro Hospitalar do Baixo Vouga, Internal Medicine Department, Aveiro, Portugal

Background: Respiratory sounds (RS) are the outcome most related to the movement of air within the tracheobronchial tree. Their characteristics are promising to easily reflect lung function in patients with chronic obstructive pulmonary disease (COPD). However, RS measurement properties, namely reliability and validity, have been poorly tested.

Aims: This study assessed the between-days reliability and validity of RS in patients with COPD.

Methods: 50 patients (36♂, 67.3±9.3y, FEV1 49.5±19.7% predicted) with stable COPD were recruited. Lung function (FEV1% predicted) was recorded with a spirometer (MicroLab 3535, CareFusion). RS at anterior and posterior right/left chest were simultaneously recorded for 20s, using air-coupled electret microphones (C417 PP, AKG), in 2 sessions conducted 5-7 days apart. Adventitious (no. of crackles and wheeze occupation rate-%wh) and normal (median frequency-F50 and maximum intensity-I_{max}) RS were processed using validated algorithms. Between-days reliability was calculated with intraclass correlation coefficient (ICC_{1,2}). Construct validity, against FEV1% predicted, was explored with Spearman's coefficient.

Results: During inspiration, no. of crackles and %wh showed excellent (ICC>0.75) and moderate-to-good (ICC>0.4) reliability at anterior and posterior chest (p<0.05). F50 and I_{max} showed moderate-to-good (ICC>0.4 and ICC>0.6, respectively) reliability at anterior and posterior right chest (p<0.05). During expiration, %wh, F50 and I_{max} were only reliable at anterior chest (ICC>0.5, ICC>0.4 and ICC>0.6, respectively; p<0.05). No. of crackles were not reliable. Significant (p<0.05) low-to-moderate (0.3<rs<0.7) correlations between FEV1% predicted and RS were found for inspiratory and expiratory no. of crackles at anterior left and posterior chest, inspiratory F50 at anterior left chest and inspiratory and expiratory I_{max} at anterior chest.

Conclusion: Inspiratory no. of crackles, recorded at posterior chest, and I_{max}, recorded at anterior chest, seem to be the most reliable and valid RS parameters to reflect lung function in patients with COPD.

Monitoring of nocturnal respiratory symptoms in COPD patients by using LEOSound lung sound monitor

Patrick Fischer¹ (patrick.fischer@ges.thm.de), Lisa Steinbrecher¹, Florian Schudt¹, Keywan Sohrabi¹, Ulrich Koehler², Andreas Weissflog³, Volker Gross¹

¹ Faculty of Health Sciences, University of Applied Sciences, Giessen, ² Department of Pneumology, Intensive Care and Sleep Medicine, University Hospital of Marburg and Giessen, Marburg, Germany

³ Thora Tech GmbH, Giessen

Background and aim: Chronic obstructive pulmonary disease (COPD) is one of most common causes of death worldwide. Typical respiratory symptoms are breathlessness, chronic cough and wheezing. Especially for night time symptoms there is a lack of awareness, which can lead to disturbed sleep and impaired daytime performance. COPD is also showing a high coincidence with obstructive sleep apnea (overlap-syndrome), which is also associated with acoustic symptoms snoring and apnea. By using objective lung sound monitoring our aim was to investigate the ability of LEOSound to monitor nocturnal respiratory symptoms like cough, respiratory rate and wheezing.

Methods: All recordings were performed by using LEOSound (Löwenstein Medical GmbH & Co. KG, Germany). LEOSound is an objective lung sound monitoring device, which contains algorithms to automatically detect coughing and wheezing.

Results: Our dataset contains recordings of 48 patients with stable COPD. Of these patients, 18 are classified into COPD stage II, 19 into COPD stage III and 11 into COPD stage IV. Most patients were male (63%) and had a mean age of 67.1 ± 7.6 years and mean packyears of 43.9 ± 20.6 .

The validation of cough and wheezing detection were performed for all 48 patients. The automated detection of cough shows a sensitivity of over 95 % and a specificity of over 80 %. Sensitivity and specificity for wheezing detection above 90 %. The automated detection of respiratory rate has a root mean square error (RMSE) ≤ 2 breaths per minute.

Conclusion: LEOSound lung sound monitor is able to record and detect cough and wheezing with high sensitivity and specificity. As a certified medical device, it offers an opportunity to record nocturnal respiratory symptoms in COPD-Patients. Other respiratory symptoms like snoring and apnea were also be recorded by using LEOSound and will be automatically detected soon.

Lung sounds intensity: is there a difference between spontaneous and target airflow?

Jácome, Cristina¹ *cristina.i.jacome@uit.no*, Aviles-Solis, Juan Carlos¹; Melbye, Hasse¹

General Practice Research Unit, Department of Community Medicine, The Arctic University of Norway.

Background: Lung sounds intensity (LSI) is a non-invasive measure to assess respiratory system that has a linear relationship with airflow. Clinicians often assess LSI asking subjects to breathe deeply with open mouth, while researchers have been using target airflows. For clinical practice, however, it would be easier to rely on lung sounds acquired without airflow measurements.

Aims: This study assessed if LSI was different when acquired with subjects breathing spontaneously and at a target airflow.

Methods: A total of 1020 recordings from 85 adults aged 40 or over were used. Lung sounds were recorded consecutively at 6 chest locations (right and left: upper posterior, lower posterior, upper anterior) using a stethoscope with a microphone in the tube. Recordings were first performed at spontaneous deep breathing and then at a target airflow of 1.5L/s. Breathing phases were manually annotated and inspiratory and expiratory LSI in the 100–2000Hz band were determined.

Results: LSI at a target airflow was found to be significantly higher than at spontaneous airflow both during inspiration (mean differences (MD) from 1.8 to 3.7dB; $p < 0.001$) and expiration (MD from 1.5 to 3.2dB; $p < 0.001$) at all locations. In inspiration, differences between right and left were only seen at upper posterior chest at spontaneous airflow ($p = 0.016$) and at lower posterior chest at target airflow ($p = 0.030$). In expiration, however, differences were seen at all locations both at spontaneous and target flows ($p < 0.034$).

Breathing phase	Location	Right		Left	
		Spontaneous	Target	Spontaneous	Target
Inspiration	Upper anterior	20.9±2.5	24.3±3.0	20.6±2.6	24.3±3.3
	Upper posterior	19.5±2.1	21.3±2.6	18.9±2.0	21.7±2.8
	Lower posterior	20.2±2.5	23.3±2.6	20.6±2.5	23.8±2.7
Expiration	Upper anterior	18.9±2.0	22.1±2.9	18.0±2.1	20.8±2.7
	Upper posterior	18.0±2.4	19.6±2.7	17.1±2.0	19.1±2.5
	Lower posterior	16.5±2.1	18.6±2.2	17±2.0	19.6±2.6

Table 1. LSI at spontaneous and target airflow (dB) (n=85).

Conclusion: LSI was higher at a target airflow of 1.5L/s than at spontaneous airflow. Despite differences in LSI were minor, its clinical significance is still unknown. This knowledge gap should direct future research.

The influence of adventitious sounds and artifacts on the frequency and intensity of normal respiratory sounds

Jácome, Cristina¹ *cristina.i.jacome@uit.no*, Aviles-Solis, Juan Carlos¹; Pasterkamp, Hans²; Melbye, Hasse¹
 General Practice Research Unit, Department of Community Medicine, The Arctic University of Norway,
²University of Manitoba.

Background and aim: Normal respiratory sounds (NRS) constitute a non-invasive way to assess lung function as they are generated by the airflow in the respiratory tract. Studies have been characterizing NRS using two main approaches: i) analyzing the complete sound file or ii) analyzing the sound file after excluding adventitious sounds/artifacts. The use of distinct approaches make comparisons between studies difficult, however, it is not known if characteristics of NRS differ significantly between the two approaches. Therefore, this study aimed to compare the differences in the spectral characteristics of NRS between the two approaches.

Methods: A total of 1020 recordings from 85 adults aged 40 or over were recorded. Lung sounds were recorded consecutively at 6 chest locations (right and left: upper posterior, lower posterior, upper anterior) using a stethoscope with a microphone in the tube. Recordings were first performed at spontaneous deep breathing and then at a target airflow of 1.5L/s. Breathing phases were manually annotated in all recordings, and then the two strategies to analyze the spectral characteristics of NRS in the 100–2000Hz band were applied. In strategy 1, without further analysis, the quartile frequencies (F25, F50 and F75) and mean intensity (Imean) were determined. In strategy 2, the quartile frequencies and mean intensity were determined after excluding adventitious respiratory sounds (crackles/wheezes) and artifacts detected with algorithms. Paired t-tests were used to analyze differences between the two strategies.

Table 1. Normal respiratory sounds at spontaneous and target airflows using two different strategies.

	Inspiration		Mean difference	p-value	Expiration		Mean difference	p-value
	Strategy 1 Mean (SD)	Strategy 2 Mean (SD)			Strategy 1 Mean (SD)	Strategy 2 Mean (SD)		
Spontaneous airflow (n=510)								
F25 (Hz)	166.9±16.0	166.5±15.9	0.39	<.001	158.2±17.5	157.2±18.3	0.93	<.001
F50 (Hz)	367.0±35.6	365.7±35.5	1.32	<.001	354.6±39.8	351.5±41.5	3.05	<.001
F75 (Hz)	664.5±44.5	662.2±44.9	2.28	<.001	656.5±46.5	651.8±50.8	4.73	<.001
Imean (dB)	20.1±2.5	20.0±2.5	0.14	<.001	17.6±2.3	17.3±2.4	0.33	<.001
Target airflow (n=510)								
F25 (Hz)	179.5±16.5	179.2±16.6	0.31	.001	169.6±18.8	168.7±19.1	0.94	<.001
F50 (Hz)	391.6±35.0	390.6±35.2	1.02	<.001	376.3±40.7	373.9±41.1	2.40	<.001
F75 (Hz)	690.3±43.3	688.7±43.3	1.62	<.001	677.1±44.6	673.1±44.3	4.03	<.001
Imean (dB)	23.1±3.1	23.0±3.1	0.14	<.001	20.0±2.9	19.6±2.9	0.35	<.001

Conclusion: Spectral characteristics are different between the two approaches, with frequency and intensity parameters achieving slightly higher values when using the first strategy. Future research should explore if these differences are clinically relevant or not.

Using machine learning to provide automatic image annotation for wildlife camera traps in the Arctic

Håvard Thom^a, John Markus Bjørndalen^a(jmb@cs.uit.no), Eivind Flittie Kleiven^b, Eeva Marjatta Soininen^b, Siw Turid Killengreen^b, Dorothee Ehrich^b, Rolf A. Ims^b, Otto Anshus^a, Alexander Horsch^a

^a Department of Computer Science, University of Tromsø - The Arctic University of Norway

^b Department of Arctic and Marine Biology, University of Tromsø - The Arctic University of Norway

Summary

The arctic tundra is considered the terrestrial biome expected to be most impacted by climate change, with temperatures projected to increase as much as 10 °C by the turn of the century. The Climate-ecological Observatory for Arctic Tundra (COAT) project monitors the climate and eco-systems using several sensor types. We report on results from projects that automate image annotations from two of the camera traps used by COAT: an artificial tunnel under the snow for capturing information about small mammals, and an open-air camera trap using bait that captures information of a range of larger sized birds and mammals. These traps currently produce over two million pictures per year.

We have developed and trained several Convolutional Neural Network (CNN) models to automate annotation of images from these camera traps. Results show that we get a high accuracy: 97.84% for tunnel traps, and 94.1% for bait traps. This exceeds previous state of the art in animal identification on camera trap images, and is at a level where we can already relieve experts from manual annotation of images.

Introduction

The arctic tundra is expected to be more challenged by climate change than any other terrestrial biome, with models projecting an average temperature increase in the Arctic as large as 10 °C by the turn of the century [1]. The Climate-ecological Observatory for Arctic Tundra (COAT) [1] [2] is a long-term research initiative for real time detection, documentation and understanding of climate impacts on terrestrial arctic ecosystems. COAT is a collaboration of several Norwegian research institutions under the umbrella of FRAM - High North Centre for Climate and Environment. The study areas include the bioclimatic extremes of the terrestrial Arctic, the low-arctic coast of the Norwegian mainland and the high-arctic Svalbard archipelago.

COAT uses several types of sensors for observations, including satellites, snow observations, meteorological observations, camera traps, and microphones. Manual annotation and analytics of images from the camera traps is by now already a demanding and time consuming task, and this burden will further increase as more traps are deployed.

This paper reports on results from using machine learning techniques on two separate sets of camera trap images: camera traps used in small mammal tunnels underneath the snow and baited camera traps above the snow. These traps are expected to capture more than two million pictures per year with the current and near future set of cameras.

Small mammal camera traps




Figure 1: Image of a tundra vole from one of the COAT camera traps under the snow.

One of the camera traps used is an artificial tunnel (see Figure 1) with a motion sensitive camera that records mammals that run through all year round [3]. These traps give brand new information on a whole community of different species (shrews, voles, lemming, least weasel and ermine) during the long arctic winter, since the traps become part of tunnel systems used by mammals in the winter. The resulting image sequences can be used to analyze animal population and community dynamics. Correct species identification is however essential for this purpose. An advantage in this regard is that the images are similar with respect to background and distance to the animal.

We trained three different Convolutional Neural Network (CNN) models and used them to predict and label images (see [4] for more details). To train the models, we had to manually annotate a dataset selected from 74,429 unlabeled camera trap images. The images were pre-processed to accommodate the algorithms we used, and to remove unnecessary information that could influence training, such as the black borders displaying image metadata.

Results show that we achieve 97.84% accuracy, 97.81% precision and 93.45% recall on a dataset with 10000 camera trap images and 11 classes. This exceeds previous state of the art in animal identification on camera trap images [5] [6] [7].



Figure 2: Image from one of the COAT open-air camera traps showing a golden eagle.

Open-air baited camera traps

These camera traps take images every five minutes, both day and night, overlooking an area with bait on the ground. Automatic annotation of these images is a harder problem than for the tunnel traps described above, as there will be more variation in weather, area around the camera, types of animals and distances to the animals in the image. Many images contain animals that are far away from the bait, making them hard to see, even by human experts. Images with animal presence will usually contain several different species, making the problem significantly harder than with single animal images. Furthermore, accumulating snow covers the bait, which partially obstructs the view of animals digging down to the bait, as seen in Feil! Fant ikke referansebildene..



Figure 3: Image from baited trap with a partially obstructed red fox surrounded by two hooded crows and one raven.

This project created a system that use three different CNN methods (Faster Region-based CNN, Single Shot Multibox Detector and You Only Look Once v2) for training and annotation, and compared the results of each method. See [8] for more details about the system.

Results from the project show that we can detect and classify animals in the images with 94.1% accuracy at 21 frames per second, exceeding the performance of previous work. Reindeer is the most challenging species to detect, as they are frequently far away from the camera. Many of the other animal classes had an accuracy close to 100%.

Conclusions

The promising results from these projects show that we are close to automating the annotation of images from the camera traps. We have not yet done a study that compares the accuracy of the automatic annotation systems to human experts that are currently annotating the images. We expect that the automatic system is close to the accuracy level of experts, as it is easy to make subtle mistakes in the software they are currently using, and humans are prone to make mistakes in tiresome and repetitive work.

We believe that the results are promising and expect that further improvements to automatic annotation of images will provide the means for scaling up the number of camera traps. This will enable observations with more spatial resolution and of larger geographic areas without the limitations currently introduced by limited human resources.

References

- [1] R. A. Ims, J. U. Jepsen, A. Stien and N. G. Yoccoz, "Science Plan for COAT: Climate-Ecological Observatory for Arctic Tundra," Fram Centre, Norway, 2013.
- [2] "COAT Web Page," [Online]. Available: <http://www.coat.no/>. [Accessed 08 2017].
- [3] E. M. Soininen, I. Jensvoll, S. T. Killengreen and R. Ims, "Under the snow: a new camera trap opens the white box of subnivean ecology.," *Remote Sensing in Ecology and Conservation*, vol. 1, no. 1, pp. 29-38, 2015.
- [4] H. Thom, "Automatic rodent identification in camera trap images using deep convolutional neural networks.," UiT - The Arctic University of Norway, Capstone Project, 2016.
- [5] X. Xu, J. Wang, R. Kays, P. A. Jansen, T. Wang and T. Huang, "Automated identification of animal species in camera trap images," *EURASIP Journal on Image and Video Processing*, vol. 2013, no. 1, p. 52, 2013.
- [6] G. Chen, T. X. Han, Z. He, R. Kays and T. Forrester, "Deep convolutional neural network based species recognition for wild animal monitoring," in *ICIP*, 2014, pp. 856-862.
- [7] A. Gomez, A. Salazar and F. Vargas, "Towards automatic wild animal monitoring: Identification of animal species in camera-trap images using very deep convolutional neural networks," vol. abs/1603.06169, 2016.
- [8] H. Thom, "Unified Detection System for Automatic, Real-Time, Accurate Animal Detection in Camera Trap Images from the Arctic Tundra," UiT - The Arctic University of Norway, Master Project, 2017.
- [9] Å. Ø. Pedersen, A. Stien, E. Soininen and R. A. Ims, "Climate-ecological observatory for arctic tundra-status 2016," *Fram Forum*, pp. 36-43, Mar 2016.

A New Computerized Approach for Lung Sounds Analysis Using Sound Signature

Omri Adler¹, Hod Gilad¹, Liran Avraham², Doron Adler² (doron_a@bat-call.com).

¹Technion – Israeli Institute of Technology, ²Bat-Call – Nesher, Israel

Background: Obstructive Lung disease is a condition where the lung airways change their shape (e.g. become narrow). These changes are associated with changes in the sounds generated within the lung during breathing. Listening to these sounds is limited by the low sensitivity of the human auditory system and the inability of the stethoscope to transfer low energy sounds.




Fig. 1: Intensity Map of Sound Signature

Aim: Suggesting a new approach to allow computerized analysis in order to overcome these obstacles.

Methods: The new method is based on Discrete Time-Frequency Distribution. It is based on Fourier transform applied to autocorrelation type function of the semi-periodical lung sound. The concept starts by sampling wide frequency lung sounds (from infra sound to audible range), identifying the active breathing periods, convert the data into its Sound Signature using the above transform, and displaying the Sound Signature (display formats are shown in the Figures).



Fig. 2: Gray Scale of Sound Signature

Results: Using this transform, we show, in this presentation, that among the properties of the proposed transform, one can find: time shift and

modulation preservation, representation of time-frequency energy distribution, and very high time-frequency resolution (limited by Heisenberg Uncertainty Principle). The new approach does not suffer from distortion of the outcome shape, while other methods, using windowing, deform the time-frequency representation (mainly by a way of low pass filtering effect). Examples of such transforms include: spectrograms, Wavelet, or Short-Time Fourier Transform. In this presentation, the Sound Signature is demonstrated on few Sound Signature examples of different pathological lung sounds, in addition examples of the Sound Signature display are presented.

Conclusion: By using the Sound Signature the lung sounds are converted from an acoustic signal into a HD digitized image. This method allows computerized classification of lung sounds using image processing classification techniques.

Acknowledgment: The authors wish to thank Prof. Noam Gavriely for his helpful review, comments and suggestions.

Use Wide Range Sound Signature in Computerized Classification of Lung Diseases

Omri Adler¹, Hod Gilad¹, Liran Avraham², Doron Adler² (doron_a@bat-call.com).

¹Technion – Israeli Institute of Technology, ²Bat-Call – Nesher, Israel

Background: Computerized classification of chest sounds is a challenging task. However, using advanced processing technologies facilitates a clinically usable medical device. Our method integrates the following: wide range pickup detector (Infrasound and audible sound), converting the data into Sound Signature images, and applying image classification Big Data algorithms.

Aim: The scope of this project is to evaluate the feasibility of implementing computerized classification to wide range lung Sound Signature.

Methods: The computerized pickup unit contain the following steps: Capturing data from the wide band pickup unit, segmentation of the breathing sequence, Sound Signature transform and displaying the data. We asked medical students to use the pickup device during regular operation of the pulmonary clinics in a few medical centers. A group of 42 volunteers participated in the study. For each volunteer we measure the lung sounds at 14 points located: six on the front chest, six on the back, and two on the chest sides. The group was divided into three: 19 Healthy volunteers, 13 COPD patients, and 10 IPF patients.

Results: The healthy Sound Signature vs. the COPD and the IPF can be separated by their patterns, as seen in the figures below:

Out of the 42 volunteers we measure, 79% of the healthy volunteers had the same Sound Signature for healthy lung sound as shown on Fig. 1, 77% of the COPD patients and 70% of the IPF patients had the corresponding Sound Signature as Presented in Fig.1.

Conclusions: The use of Infrasound as part of the lung Sound Signature enhances the classification of chest sounds and may facilitate the calculation of probability-of-existence of certain disease and screening lung conditions.




Fig. 1: Example of three Sound Signatures. Left – Healthy, Center – patient with COPD, Right – Patient with IPF.

Acknowledgment - The authors wish to thank Prof. Noam Gavriely for his helpful review, comments and suggestions.

Detection of Wheezes and Breathing Phases Using Deep Convolutional Neural Networks

Johan Ravn (jonna12_@hotmail.com)¹, Juan Carlos Aviles Solis², Morten Grønnesby³, Hasse Melbye², Lars Ailo Bongo⁴

¹ Medsensio AS, N-9019 Tromsø, Norway, ² General Practice Research Unit in Tromsø, Department of Community Medicine, UiT The Arctic University of Norway, N-9037 Tromsø, Norway. ³ Department of Medical Biology, UiT The Arctic University of Norway, N-9037 Tromsø, Norway. ⁴ Department of Computer Science, UiT The Arctic University of Norway, N-9037 Tromsø, Norway.

Introduction: Detection and interpretation of lung sounds is difficult, so inexperienced health personnel classify lung sounds differently [1]. Integrating an automatic system for detection and classifying lung sounds into digital stethoscopes could help interpret lung sounds. Previous approaches for detection of wheezes have used small datasets and machine learning methods such as Gaussian mixture models [2], artificial neural networks [3] and support vector machine [4].

We use a variant of artificial neural networks called deep learning. The recent improvements in deep learning algorithms, data availability, and computational power have allowed deep learning to be used in fields such as self-driving cars, image analysis, and speech recognition. Also, in medical fields, large image datasets have been used to train deep learning models to automatically classify symptoms and diseases [5]. However, to our knowledge, deep learning for classification has not been applied to lung sounds.

Methods: Deep learning typically requires a bigger dataset than most other machine learning approaches. The Tromsø Study 7 has collected a large lung sound dataset of which we use 8784 unique recordings of lung sounds. We have developed deep learning algorithms and trained them using this dataset. The algorithms detect wheezes and breathing phases in lung sounds. We have used the algorithm to build a lung sound classification service that can be used by various applications such as a web based lung auscultation learning tool for students.

Our approach uses state-of-the-art deep convolutional neural networks (ConvNets), to automatically detect breathing phases and wheezes in lung sound recordings. Using ConvNets, we can utilize large datasets without manual feature engineering. Related work has shown that deep ConvNets can outperform previous methods for acoustic recognition on tasks such as speech recognition and acoustic event detection.

We evaluate our wheeze detection system using recall and precision. We also calculate the mean class precision and recall across both classes. For the breathing phase detection system, we use mean average precision (mAP). This metric is often used to evaluate results from a document retrieval system, but also for object detection systems.

Table 1: Wheeze detection.

Class	Precision (%)	Recall (%)
Normal	79	98
Wheeze	96	73
Mean class	89	85,5

Table 2: Breathing phase detection.

Class	AP (%)
Inspiration	89
Expiration	88
mAP	88

Results: Our results (Table 1 and Table 2) show that we have successfully implemented wheeze detection with wheeze recall (sensitivity) of 73%, and normal recall (specificity) of 98%. We have also implemented a breathing phase detection with an average precision of 89% for the inspiration phase and 88% for the expiration phase.

Figure 1. Screenshot from our lung auscultation prototype used to detect wheezes.





Figure 4. Screenshot from our lung auscultation prototype used to detect breathing phases.

Spectrogram with annotations



Conclusion and future work: Our approach successfully detects wheezes and breathing phases. Further research is necessary to determine how we can integrate our approach into commercial applications such as a learning tool, a digital stethoscope or a home monitoring device. Encouraged by these results, we have started a company Medsensio AS, which have received initial funding from the Research Council of Norway. Our main goal will be to use novel methods and developments in computer science for medical data analysis. We will start by using our approach in a lung auscultation teaching tool and integration with smart stethoscopes.

We are currently investigating the clinical and research aspects of lung sounds, such as how lung sounds are used for early screening and the relationship with diseases such as COPD and asthma. Furthermore, we plan to validate our approach on other datasets recorded in different environments and with different equipment.

Bibliography

- [1] J. C. Aviles-Solis, P. Halvorsen and H. Melbye, "Inter-observer variation in categorizing lung sounds," in *The International Lung Sound Association 42nd Annual Conference*, St. Petersburg, 2015.
- [2] M. Bahoura and C. Pelletier, "Respiratory Sounds Classification using Gaussian Mixture Models," in *Electrical and Computer Engineering*, 2004.
- [3] B.-S. Lin, H.-D. Wu and S.-J. Chen, "Automatic Wheezing Detection Based on Signal Processing of Spectrogram and Back-Propagation Neural Network," *Journal of Healthcare Engineering*, vol. 6, no. 4, pp. 649-672, 2015.
- [4] D. Chamberlain, R. Kodgule, D. Ganelin, V. Miglani and R. R. Fletcher, "Application of Semi-Supervised Deep Learning to Lung Sound Analysis," in *38th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC)*, 2016.
- [5] V. Gulshan, L. Peng, M. Coram and et al, "Development and Validation of Deep Learning Algorithm for Detection of Diabetic Retinopathy in Retinal Fundus Photographs," *JAMA*, vol. 316, no. 22, pp. 2402-2410, 2016.

Machine Learning Based Crackle Detection

Morten Grønnesby (morten.gronnesby@uit.no)¹, Juan Carlos Aviles Solis,² Einar Holsbø,³ Hasse Melbye,² Lars Ailo Bongo³

¹ Department of Medical Biology, UiT The Arctic University of Norway, N-9037 Tromsø, Norway.

² General Practice Research Unit in Tromsø, Department of Community Medicine, UiT The Arctic University of Norway, N-9037 Tromsø, Norway.

³ Department of Computer Science, UiT The Arctic University of Norway, N-9037 Tromsø, Norway.

Background: In recent years, many innovative solutions for recording and viewing sounds from an electronic stethoscope have become available. However, to fully utilize such smart stethoscopes, there is a need for an automated approach for detecting abnormal lung sounds.

Aim: To create an algorithm for automated detection of crackles in lung sound recordings.

Methods: We developed a machine learning based approach for detecting crackles in lung sounds recorded using a stethoscope with a microphone in the tube in the 7th Tromsø Study (a health survey). We trained and evaluated the machine learning model using 209 files with crackles classified by experts. We use features extracted from small windows in audio files. We evaluated several feature extraction methods and classifiers.

Results: We got the best performance using four features from the time domain and one from the spectrum domain, classified by a SVM with a Radial Basis Function Kernel. With this approach, we obtained a precision of 86% and recall of 84% for classifying a crackle in a window. This is more accurate than found in human classifiers in studies of health personnel. The classifier is very fast.

Conclusion: Our approach detects and visualizes individual crackles in recorded audio files. It is accurate, fast, and has low computational resource requirements. The approach is therefore well suited for development of tools that can help health personnel to detect and understand the implication of crackles. Further research is necessary to demonstrate the use of such tools in the clinical setting.

Comparison of Various Machine Learning Algorithms on Adventitious Pulmonary Sound Classification

Sezer Ulukaya^{1,2}, Gorkem Serbes³ and Yasemin P. Kahya (kahya@boun.edu.tr)¹

¹Dept. of Electrical and Electronics Engineering, Boğaziçi University, Istanbul, Turkey

²Dept. of Electrical and Electronics Engineering, Trakya University, Edirne, Turkey

³Dept. of Biomedical Engineering, Yildiz Technical University, Istanbul, Turkey

{sezer.ulukaya,kahya}@boun.edu.tr, {gserbes}@yildiz.edu.tr

Background: Time, location and frequency of adventitious pulmonary sounds may help to track signs of different pulmonary diseases. In literature, various techniques are proposed to classify pulmonary sounds using machine learning algorithms.

Aim: In this study, using various machine learning algorithms pulmonary sounds are classified into normal vs. adventitious groups. Moreover, continuous pulmonary sounds are further classified into mono/polyphonic wheeze types.

Methods: Non-dyadic wavelet transform is used to extract raw features from pulmonary sounds by comparing Fourier transform and dyadic discrete wavelet transform methods. Machine learning algorithms such as k Nearest Neighbour, naive Bayes and support vector machines are used to compare classifier's performances.

Results: The experimental results showed that non-dyadic wavelet transform performs better classification accuracies when compared to Fourier based methods with 5-10 % higher correct classification rates both in normal-adventitious and wheeze type classification problems. Support vector machine classifier with radial basis function performs the best correct classification rates when raw features are fed into the classifiers. Performance of various parameters are provided to demonstrate optimum parameter set also.

Conclusion: Wavelet based technique provides better time frequency resolution than Fourier transform and dyadic discrete wavelet transform methods, however in case of wheeze type classification an adaptive wavelet based method is needed. Support vector machine classifier when compared to other classifiers provides better results.

Acknowledgements: The work is supported by Boğaziçi University Research Fund under grant number 16A02D2. The work of S. Ulukaya is supported by the Ph.D. scholarship (2211) from Turkish Scientific Technical Research Council (TUBITAK).

Cardiac Response to Respiration in Deep Inspiration followed by Breath Holding

Sadamu Ishikawa M.D. (sadaishi@massmed.org) and Peter La Camera M.D.

Pulmonary and Critical Care, Steward St Elizabeth's Medical Center

Dept. of Medicine Tufts University School of Medicine, Boston, Ma. U.S.A.

US -1-617-789-2563

Background: Heart rate becomes slower on inspiration. When one takes deeper breathe, as more negative pressure is generated within the chest which leads to more blood returning to left ventricle, then a larger volume of blood must be pumped out, hence a delay of heart beats on inspiration. We have been studying this phenomenon for past 4 years and reported at annual ILSA meetings on healthy nonsmoking subjects, smokers, and patients with COPD.

Methods: This year we investigated Effects of Breath Holdings in 10 Healthy Nonsmoking Individuals when they come to my office for routine visit with no acute respiratory illnesses. We used 2 channel ECG and lung sounds simultaneously recording ECG and tracheal sounds at sitting position.

Results: 2 QRS intervals became 41 msec on Deep Inspiration comparing to Expiration Period of 30 msec. average of all 10 subjects, mean heart rate reduction from 76 to 59 /min.

While deep inspiration followed by breath holding, heart rate reduction occurred comparing to expiration period. After change of 2QRS intervals following deep inspiration (became longer, slowing heart rate) the 2 QRS intervals remained essentially the same during breath holding period.

Conclusion: We conclude that deep inspiration with breath holding is more advantageous for slowing heart beat comparing to just deep inspiration.

Auscultatory phenomena in patients with partially controlled bronchial asthma with intrapulmonary auscultation

Sergei Glotov, Yuriy Petrov (petrov.yuriy1992@gmail.com), Vladimir Abrasimov.

Department of postgraduate education in Internal Medicine, Ryazan medical university, Ryazan, Russian Federation

Background: One of the most common chronic diseases is bronchial asthma, and in recent years, the incidence of it is only increasing. This represents a serious medico-social and economic problem, especially against the backdrop of untimely and inadequate diagnosis of the disease.

Materials and methods: During the traditional bronchoscopy (Karl Storz Tele Pack X) is performed simultaneously with video bronchoscope intrapulmonary registration acoustic phenomena specifically designed to work with the bronchoscope microphone. Simultaneously recording lung sounds produced by the electronic stethoscope (3M Littmann Electronic Stethoscope 3200) for further analysis and comparison data

Results: By recording in 20 patients with partially controlled asthma (GINA 2014). The main characteristics, which evaluated abnormal noises were a range of received frequencies and duration of which depend on the sensitivity of the sensor and the proximity to the site of occurrence of wheezing. This was traced frequency correlation on the cause of wheezing. Thus, the average duration of wheezing during intrapulmonary auscultation was about 250 ms, the prevailing average rate of wheezing 400 Hz, the bass - 200 Hz ($p < 0,05$). Frequency range wheezing - 80-1600 Hz ($p < 0,05$). The highest rate was observed in the presence of mucus and, especially, the spasm, which were recorded using the received video and flutter to give more bass version of wheezing. When registering the sounds on the body surface average duration was 100 ms, the prevailing average frequency - 150 Hz, and the range was only 350-950 Hz ($p < 0,05$). A clear correlation when it was received.

Conclusion: Intra-pulmonary auscultation can improve the quality of the diagnostic process by obtaining a wider range of diagnostic information, will help to better understand the nature of pathological processes occurring in the lower respiratory tract and will positively affect the effectiveness of examination and treatment of the patient.

Lung sounds of patients with interstitial pneumonia in left and right lateral decubitus position - Effect of gravity on lung sounds

Michiko Tsuchiya (rakuwadr015@rakuwadr.com), Yukio Nagasaka, Chikara Sakaguchi, Hitomi Ajimizu, Yosuke Nakanishi, Noboru Morikawa, Kiyoshi Ninomiya

Department of Pulmonary Medicine, Rakuwakai Otowa Hospital

Background: We reported that lung sounds of interstitial pneumonia (IP) were influenced by body position, and lung sound intensity (LSI) was greater in the dependent position. As most of the sound power comprised of crackles, this observation means that crackles in IP is louder in the dependent lung. We also found that LSI was not symmetrical; LSI was greater on the left than on the right and postural changes of LSI were greater on the right than on the left. This study aims to clarify the effects of lung position on LSI, i.e., dependent side or upside, when the subjects are in right and left lateral decubitus positions.

Methods: We recorded lung sounds on both the right and the left lung bases in left and right lateral decubitus position in seventeen patients with IP. Lung sounds were recorded for ten seconds by sound spectrometer, LSA2012 (Kenz-Medico). We averaged the sound intensity of whole ten seconds and compared the averaged LSI in the two lateral decubitus positions.

Results: In left lateral decubitus position, LSI on the left lung (dependent lung) was significantly greater than that on the right lung (upside lung). In right lateral decubitus position, there was no significant difference in LSI between the left lung (upside lung) and the right lung (dependent lung). The postural change of LSI between upside and dependent side was greater on the right lung than on the left lung.

Conclusions: The LSI of IP, i.e., loudness of the crackles was greater when the lung was in the dependent side than in the upside. The postural change of LSI was greater on the right lung than the left lung.




Figure: Sound spectrogram of two cases with IP. The lung sounds were recorded on bilateral lung bases in right lateral and left lateral decubitus position.

The crackles are recognized as bunch of light blue vertical lines. In both cases, crackles are more prominent in the dependent side than in the upside.

In case 1, crackles in upside lung disappear. In case 2, attenuation of crackles in left lung is slight in the right lateral decubitus position.

Table: The lung sound intensity (dBm) in each frequency range

Body position	Sound frequency	Right lung Mean (SD)	Left lung Mean (SD)
Right Lateral Decubitus	Low	5.0 (10.7)	8.1 (10.2)
	Middle	-13.5 (12.8)	-11.2 (11.6)
	High	-24.8 (12.2)	-26.5 (14.0)
Left Lateral Decubitus	Low	-4.9 (11.9)	13.5 (8.6)
	Middle	-25.3 (10.5)	-3.7 (9.3)
	High	-39.7 (10.0)	-16.0 (11.7)

Comparing lung sounds intensity between sitting, supine and prone position in patients with interstitial pneumonia

Michiko Tsuchiya (rakuwadr015@rakuwadr.com), Yukio Nagasaka, Chikara Sakaguchi, Hitomi Ajimizu, Yosuke Nakanishi, Noboru Morikawa, Kiyoshi Ninomiya

Department of Pulmonary Medicine, Rakuwakai Otowa Hospital

Background: Auscultation of patients with stable interstitial pneumonia (IP) are usually done in the sitting position, whereas in critically ill conditions, patients often lie in the supine position. Last year we reported that lung sounds of IP were influenced by body position in eight patients with IP. Most of the sound power was comprised of crackles. These findings suggest that the loudness of crackles in IP was influenced by body position. This year, we tried to assure these observations by increasing the number of the subjects.

Methods: We analyzed lung sounds of seventeen patients with IP by a sound spectrometer, LSA2012 (Kenz-Medico). Two acoustic sensors were placed on the left and the right lung bases. The lung sounds were recorded in sitting, supine and prone positions. We recorded the lung sounds for ten seconds and averaged sound intensity of whole recorded sounds. We measured the lung sound intensity (LSI) of three frequency ranges, low (L: 200-390Hz), middle (M: 400-790Hz), and high (H: 800-1600Hz), and compared the averaged LSI of each frequency ranges in each body positions.

Results : The LSI was greater on the left than the right in all body positions. The difference of LSI between the left and the right lung was the greatest in prone and the smallest in supine position, although there were some individual variations as shown in the figure. There were significant postural changes of LSI on the right but not on the left. Statistically significant difference was observed between prone and supine, and prone and sitting position on the right. However, no significant postural change of LSI was observed on the left. (Table)

Conclusions: LSI of IP was greater on the left than on the right, suggesting that the loudness of crackles were louder on the left than on the right in all body positions. The postural changes of LSI were significant on the right but not on the left. These observations should be considered in auscultation and the lung sound study of IP.




Figure. Sound-spectrogram of two cases of IP.

In both cases, crackles are more prominent in supine position. The lung sounds are more loud on the left than on the right. The effects of body position on LSI are greater in case 1 than in case 2. The differences of LSI between the right and the left are greater in case 2 than in case 1.

Body position	Sound frequency	Right lung Mean (SD)	Left lung Mean (SD)
Sitting	Low	1.0 (8.5)	8.3 (9.1)
	Middle	-16.9 (9.0)	-9.7 (14.0)
	High	-26.2 (9.4)	-19.5 (16.3)
Supine	Low	1.3 (9.4)	6.5 (12.8)
	Middle	-11.2 (9.4)	-7.2 (12.1)
	High	-24.4 (10.4)	-18.3 (13.7)
Prone	Low	-4.6 (10.3)	9.1 (9.1)
	Middle	-24.2 (13.3)	-10.6 (12.0)
	High	-37.6 (13.3)	-23.8 (13.9)

Table. The lung sound intensity (dBm) in each frequency range

Lung sound changes associated with pneumothorax

Mansy HA (PhD) hansen.mansy@ucf.com, Balk R (MD), Sandler RH (MD)

University of Central Florida Orlando, FL 32816, and Rush University medical Center,
Chicago, IL 60612

Pneumothorax (PTX) is an abnormal accumulation of air in the plural space that can be life threatening.

Purpose: The purpose of the study is to investigate the changes in lung sounds associated with the occurrence of pneumothorax.

Methods: Patients undergoing video assisted thoracic surgery (VATS) participated in the study after informed consent. Subjects were sedated and mechanically ventilated then one lung was collapsed as a normal part of the surgical procedures. Sounds were recorded at chest surface with an electronic stethoscope placed at the mid axillary line at the level of the 5th intercostal space while subjects were in the decubitus position. Measurements were repeated for the control and pneumothorax states in each patient.

Results and conclusions: A drop in the acoustic energy with PTX was observed over the affected hemithorax. This drop may be because the collapsed lung and airways interfered with sound transmission in the affected side and/or due to diverting airflow and acoustic energy to the unaffected side. These results suggest that a simple acoustic method may be used to distinguish PTX from the control state.

Crackles and Rhonchi with Postural Dependence in a 9-year-old Girl with Severe Mycoplasma Pneumonia

Masato Takase, MD, PhD (mtakase@nms.ac.jp)

Department of Pediatrics, Nippon Medical School Tama-Nagayama Hospital, Tokyo, Japan

Background: *Mycoplasma pneumoniae* (Mp) is the most common cause of pneumonia in school age children. In general, there are few physical signs such as crackles and wheezes, despite general symptoms and problems with the respiratory system. Usually, mycoplasma pneumoniae is successfully treated with macrolides or tetracyclines. However, there are a few unusual cases which do not respond to these antibiotics and require the systemic use of corticosteroids.

Case: A 9-year-old girl was admitted to our hospital for high fever and cough lasting seven days. She had been taking clarithromycin (CAM) since the 3rd day of the illness, but symptoms did not improve. On physical examination, some crackles were noticed, and a chest X-ray revealed bilateral diffuse patchy infiltrates. Her SpO₂ was 95% breathing room air and oxygen was administered through a nasal cannula. Mycoplasma pneumoniae was suspected, and intravenous sulbactam/ampicillin and oral minocycline were administered. However, her symptom worsened, and crackles and rhonchi increased dramatically. The recordings of the lung sounds confirmed coarse crackles and expiratory rhonchi which increased markedly on the dependent side of lateral decubitus position. On the 4th day of admission, prednisolone 1mg/kg/day was introduced, and the symptoms dramatically improved thereafter. The diagnosis of a mycoplasma pneumoniae was confirmed with the increased serum antibody titer (20480x) for Mp.

Conclusion: Postural dependence of crackles was said to be a characteristic of the fine crackles. However, as shown in this case, coarse crackles could have the similar characteristic in some occasion.

Rumbling rhonchi in bronchial asthma: Their relation to airway inflammation

Yukio Nagasaka (rakuwadr1127@rakuwadr.com), Michiko Tsuchiya, Youske Nakanishi, Chikara Sakaguchi,

Hitomi Ajimizu, Noboru Morikawa, Kiyoshi Ninomiya

Department of Pulmonary Medicine, Rakuwakai Otowa Hospital

Introduction: The rumbling rhonchi are low-pitched continuous rales and imply retained secretion in bronchi, implying presence of airway inflammation. (35th ILSA2010, 41st ILSA 2016, Allergol Int. 2012; 61:353) We examined parameters of airway inflammation in rather stable asthmatic patients, with or without rumbling rhonchi.

Methods: We examined pulmonary function, FENO (Fraction of exhaled nitric oxide) and peripheral blood eosinophils as parameters of airway inflammation in 148 non-smoking stable asthmatic patients (Age: 63 ± 16, M/F: 59/79) when they had no acute symptoms. Non-smoking subjects included 43 ex-smokers and 105 never smokers. Ex-smokers were involved in this study when they did not smoke for more than three years.

The loudness of the rumbling rhonchi was classified into three grades, Grade 0 (no rumbling rhonchi), Grade 1 (present, but faintly audible) and Grade 2 (present and loudly audible). There were 47 patients (10 ex-smokers and 37 never smokers) in Grade 0, 61 patients (16 ex-smokers and 45 never smokers) in Grade 1 and 40 patients (17 ex-smokers and 23 never smokers) in Grade 2.

Results: FENO was significantly higher in Grade 2 than in Grade 0 ($p < 0.01$) and Grade 1 ($p < 0.03$). There was no significant difference of FENO between Grade 0 and Grade 1 ($p = 0.23$). There was no significant difference of pulmonary function among Grade 0, Grade 1 and Grade 2. There was no significant difference of pulmonary function among Grade 0, Grade 1 and Grade 2, although peripheral blood eosinophil appeared to be higher in Grade 2.

Conclusions: We conclude that rumbling rhonchi suggest airway inflammation as shown in increase of FENO. However, presence of rumbling rhonchi did not suggest decreased pulmonary function nor eosinophilia in the peripheral blood.

Table: FENO, %FEV1/FVC and peripheral blood eosinophil (number and percentage) with the grade of rumbling rhonchi

Rumble Grade	FENO (mean +/-SD)	%FEV1/FVC (mean +/-SD)	Eosinophils /cmm (mean +/-SD)	Eosinophils (%) (mean +/-SD)
0 (no) N=47	23.4 +/- 32.7	73.5 +/- 29.1	227 +/- 479	3.6 +/- 6.4
1 (faint) N=71	30.1 +/- 31.8	79.8 +/- 11.1	199 +/- 157.2	3.7 +/- 2.8
2 (loud) N=40	55.4 +/- 67.9	73.7 +/- 11.4	509 +/- 961	5.9 +/- 8.6

Rumbling rhonchi and bronchial inflammation in patients with bronchial asthma

Yukio Nagasaka (rakuwadr1127@rakuwadr.com), Michiko Tsuchiya, Youske Nakanishi, Chikara Sakaguchi, Hitomi Ajimizu, Seiya Nishiyama, Yasuyuki Hayashi, Noboru Morikawa

Department of Pulmonary Medicine, Rakuwakai Otowa Hospital

Introduction: The rumbling rhonchi have non-sinusoidal wave in their time-expanded wave form and imply retained secretion in bronchi, thus supposed to reflect an aspect of airway inflammation. (35th ILSA2010, *Allergol Int.* 2012; 61:353) As there have been no previous studies on the clinical implication of presence of rumbling rhonchi in patients with bronchial asthma, we examined pulmonary function and FENO (fraction of expired nitric oxide) as parameters of airway inflammation in rather stable asthmatic patients, with or without rumbling rhonchi.

Methods: We studied eighty cases of non-smoking adult asthmatics (Age: 62.9 ± 15.62 , M/F: 36/31) when they had no acute symptoms. All of the ex-smokers involved in this study quitted smoking for more than three years. After careful auscultation of the lung, the patients were examined pulmonary function and FENO (fraction of expired nitric oxide). The loudness of the rumbling rhonchi was classified into three grades, Grade 0 (no rumbling rhonchi), Grade 1 (present, but faintly audible) and Grade 2 (present and loudly audible). These grades were determined by the loudness of the rumbling rhonchi irrespective of their extent.

Results: There were 18 patients (three ex-smokers and 15 never smokers) in Grade 0, 37 patients (11 ex-smokers and 26 never smokers) in Grade 1 and 25 patients (11 ex-smokers and 14 never smokers) in Grade 2. Inspiratory or expiratory short wheezes were heard in seven patients in Grade 0, three patients in Grade 1 and two patients in Grade 2. These wheezes were heard in small area, less than one quarter of anterior or posterior chest.

FENO was significantly higher in Grade 2 than in Grade 0 ($p < 0.05$) and Grade 1 ($p < 0.05$). There was no significant difference of FENO between Grade 0 and Grade 1 ($p = 0.23$). There was no significant difference of pulmonary function among Grade 0, Grade 1 and Grade 2. (Table)

Conclusions: Rumbling rhonchi reflected airway inflammation as shown in increase of FENO. However, the grade of rumbling rhonchi did not correlate with the pulmonary functions or presence of short wheezes.

Rumbling rhonchi were heard more often in patients who had history of smoking in the past. Thus, the presence of rumbling rhonchi was assumed to reflect eosinophilic airway inflammation and also chronic airway inflammation caused by smoking of more than years ago.

Table

Rumbling rhonchi	FENO (ppb)	FEV1/FVC (%)	FEF75*/predFEF75 (%)
Grade 0 (n=18)	23.6 ± 12.78	69.8 ± 12.50	53.9 ± 28.52
Grade 1 (n=37)	32.2 ± 19.20	78.1 ± 12.27	81.6 ± 47.43
Grade 2 (n=25)	63.3 ± 93.09	72.4 ± 24.29	62.0 ± 36.64

*FEF75: expiratory flow rate at 75% of FVC (forced vital capacity).

Lung sounds of a patient with bronchiolitis obliterans: comparison with pulmonary emphysema

Hiroshi Nakano (nakano_fukuoka@yahoo.co.jp), Tomoe Nishihara, Akiko Ishimatsu, Tomoaki Iwanaga
Fukuoka National Hospital, Fukuoka, JAPAN


Background: We have reported enhanced high frequency component in lung sounds of pulmonary emphysema (PE) (Ishimatsu: Intern Med 2015; 54:1183-1191). Both PE and bronchiolitis obliterans (BO) exhibit severe airflow obstruction, while underlying morphological changes are quite different between them. Thus, comparison of lung sounds between PE and BO may be important to understand the mechanisms for lung sound characteristics of obstructive lung diseases.

Methods: Subjects were a BO patient (Case report by Nishihara: Chest 2017; 151(3): e57-e62), 20 COPD patients, and 20 normal subjects. Tracheal sounds and lung sounds at six sites (R&L-upper, R&L-middle, R&L-lower) were analyzed using FFT. Sonogram and power spectra were compared.

Results


1. Adventitious sounds:

In the BO case, coarse crackles were found at the very beginning of inspiration (Fig) over all the lung fields. In the PE cases, crackles were identified in 13 out of 20 cases, only at lung bases, and not confined to the beginning of inspiration.



2. Basic breath sounds: Both BO and PE cases had greater breath sound intensity at a given airflow in both inspiratory and expiratory phases as compared with normal subjects (Fig).

Coherence analyses indicated that the origin of basic breath sounds in the BO case was located at a peripheral region not only for inspiratory but also expiratory phase, which was different from PE cases.



Conclusion: Patients with airway obstruction often exhibit breath sounds with enhanced high frequency components. The mechanism underlying this phenomenon may be different between airway diseases and parenchymal diseases.

Sub-classification of wheezes and crackles is not helpful when lung auscultation is used for detecting decreased lung function in adults

Juan Carlos Aviles Solis¹, Cristina Jácome¹, Hans Pasterkamp², Hasse Melbye¹ (hasse.melbye@uit.no)

General Practice Research Unit, Department of Community Medicine, UIT The Arctic University of Norway,
²University of Manitoba, Canada.

Background: Pulmonary auscultation is frequently a part of physical examination during health screening, with crackles and wheezes regarded as valuable, but not very specific, signs of lung disease. In the Tromsø study (2015-16) we had the opportunity to record lung sounds in a random sub-sample of 21,083 adults aged 40 years or older. The aim of the present analysis is to describe the associations between adventitious lung sounds and reduced lung function as measured by spirometry and pulse oximetry, and to assess whether subcategories of crackles and wheezes differ in their value to detect reduced lung function.

Methods: 15 seconds of lung sounds were recorded at 6 positions on the chest in 4,033 participants. The participants were asked to breathe deeply with an open mouth. The recordings were classified for the presence of crackles or wheezes (including rhonchi) by two independent observers. Cases of disagreement were discussed with a third observer, and agreement was obtained by consensus. All recordings with crackles or wheezes thus classified underwent a second evaluation by two pairs of observers. Again, crackles and wheezes were classified as present or absent, with an additional class of "uncertain". It was also possible to classify the recordings as normal. Crackles (present or uncertain) were further sub-classified as single or multiple and as coarse or fine. Wheezes were sub-classified as short, ordinary or long, and into wheezes or rhonchi, where musical low-pitched wheezes were to be classified as wheezes. When both observers in a pair classified a recording as normal or felt that the presence of crackles or wheezes was uncertain, the final classification was "normal". When both observers in a pair classified crackles or wheezes to be present, or one felt the sound was present and the other was uncertain, the sound was classified as present. When one observer found a recording normal and the other felt that crackles or wheezes were present, the recording was discussed between all four observers. If at least three out of the four observers found crackles or wheezes to be present, this was accepted as the final classification. Spirometry was carried out according to ATS/ERS guidelines using Vmax Encore spirometer. Global Lung Function Initiative reference values were used. Oxygen saturation was measured with Onyx II pulse oximeter, and the highest SpO₂ after three measurements was used. FEV₁/FVC less than the lower limit of normal (LLN), FEV₁% ≤ 70 predicted, and SpO₂ ≤ 95% were used as outcome variables. We also had information on smoking habit, and the participants answered the Modified Medical Research Council (mMRC) questionnaire on shortness of breath (score 0-4).

Results: Among the 4,033 participants with lung sounds recorded, the median age was 65 years (range 40-84), and 53.5% were women. Spirometry was available from 3,799, and pulse oximetry from 3,872 participants. Crackles were found in 532 participants (13.2%),

with purely inspiratory crackles in 495. Wheezes were found in 729 (18.1%). FEV1/FVC less than LLN was found in 10.2%, FEV1% \leq 70 in 5.2% and SpO2 \leq 95% in 4.7%. Crackles were significantly associated with all three outcome variables, and most strongly with decreased oxygen saturation (OR=2.7). Wheezes were only associated with the spirometric outcomes, and most strongly with FEV1% \leq 70 (OR=1.7). Current and previous smoking and shortness of breath (mMRC \geq 2) were stronger predictors of all outcome variables, with ORs between 2.3 to 7.3 (current smoking for FEV1% \leq 70). Sub-classification of inspiratory crackles showed a somewhat higher OR of multiple vs. single crackles regarding decreased oxygen saturation (figure 1) but the difference was not statistically significant. Even less difference was found between fine and coarse crackles. For wheezes (inspiratory or expiratory) there was a somewhat higher OR of ordinary and long vs. short wheezes regarding FEV1 % \leq 70 (figure 2), but the difference was not statistically significant. Even less difference was found between wheezes and rhonchi. There was very low agreement between the observers for all sub-classifications (figure 1 and 2).

Conclusion: Crackles and wheezes are significant predictors of reduced lung function in the general population, but weaker predictors than smoking habit and shortness of breath. Subcategories of crackles and wheezes were not significantly stronger predictors of reduced lung function. Since the agreement between observers on subcategories is very low, it is recommendable to not pay attention to sub-classifying when using lung auscultation as a tool in population screening of adults.

Fig 1. Odds ratio of inspiratory crackles, including sub-categories, for SpO2 \leq 95%





Fig 2. Odds ratio for wheezes, including sub-categories, for FEV1% predicted \leq 70



Can normal respiratory sounds differentiate COPD grades?

Miranda, Sara^{1,2} Jácome, Cristina^{1,3} Machado, Ana^{1,2} Melro, Hélder^{1,2} Simão, Paula⁴ Marques, Alda^{1,2}
(amarques@ua.pt)

¹ Lab3R – Respiratory Research and Rehabilitation Laboratory, School of Health Sciences, University of Aveiro, Aveiro, Portugal, ² iBiMED – Institute for Research in Biomedicine, University of Aveiro, Aveiro, Portugal, ³ Department of Community Medicine, UIT the Arctic University of Norway, ⁴ Pulmonology Department, Hospital Pedro Hispano, Unidade Local de Saúde de Matosinhos, Matosinhos, Porto, Portugal

Background: Normal respiratory sounds (NRS) are a reliable and useful marker for the diagnosis and monitoring of chronic obstructive pulmonary disease (COPD) and are valuable for early detecting pathological events through the course of the disease. However, little is yet known about its characteristics in different severity grades of the disease.

Aims: This study explored the characteristics of NRS across COPD grades. **Methods:** 66 patients with COPD were recruited from routine pulmonology appointments. Lung function data was used to categorize patients according to GOLD severity classification into two groups, either mild-to-moderate (n=30; 64.9±13.7yrs; 60%♂; FEV1 72±18% pred) or severe-to-very-severe (n=36; 67.5±7.5yrs; 75%♂; FEV1 33±11% pred) grades. NRS were recorded with a microphone for 20 seconds on the posterior left chest, at spontaneous airflow. NRS frequency and intensity were analyzed with validated algorithms within the 100-300 Hz band. Mann–Whitney U tests were used.

Results: The groups were homogeneous in terms of age and gender. No significant differences were found between the two groups in NRS during both inspiration and expiration in the median frequency and mean intensity (Table 1).

Conclusion: NRS seem to be similar between different COPD severity grades in stable patients. Further studies with larger samples are needed to enhance our understanding on the respiratory acoustics during the course of COPD.

Table 1. Frequency (Hz) and mean intensity (dB), posterior left chest, 100-300 Hz, by disease severity (n=66)

	Inspiration			Expiration		
	GOLD 1 and 2 (n=30)	GOLD 3 and 4 (n=36)	p-value	GOLD 1 and 2 (n=30)	GOLD 3 and 4 (n=36)	p-value
F25	36.2 [26.2 - 40.0]	34.0 [25.6 - 37.7]	p=0.776	25.4 [21.3 - 33.4]	26.9 [21.7 - 36.0]	p=0.333
F50	80.4 [60.9 - 85.5]	76.3 [57.9 - 85.2]	p=0.285	58.5 [48.4 - 77.6]	61.0 [52.0 - 80.0]	p=0.285
F75	130.3 [109.1 - 139.0]	125.3 [104.3 - 137.8]	p=0.842	105.6 [88.1 - 129.2]	109.4 [93.5 - 131.5]	p=0.341
Mean intensity	16.1 [9.9 - 25.9]	17.7 [11.2 - 32.7]	p=0.455	9.3 [7.2 - 19.2]	13.2 [8.7 - 23.8]	p=0.094

Is Auscultation Still Useful or is it Just Nostalgia?

Steve Kraman, M.D. (sskram01@uky.edu)

Professor, Division of Pulmonary, Critical Care and Sleep Medicine, Department of Internal Medicine, University of Kentucky, Lexington, Kentucky, USA

Not surprisingly, I have an opinion on this matter. In fact, I believe that auscultation of the chest is not only as important as it ever was, but is more important now than in the past. The reason for this is simply that we can now do so much more for diseases that are discovered during the history and physical examination. This makes rapid diagnosis more critical than it has ever been. For example, if in the 1850s, a middle-age man suddenly complained of indigestion, whether it was truly indigestion or a myocardial infarction hardly mattered as there was little that could realistically be done for him at the time. He would either recover or not. Now, he would be rushed to an emergency room where something as simple as listening to his chest and discovering extrasystoles or a new mitral regurgitation murmur would point the physicians in the right direction and (following an ECG) he would be in the Cath Lab within 90 minutes. Distinguishing heartburn from a heart attack is of critical importance.

Similar arguments can be made for pneumonia. Before the antibiotic era, a physician could only apply a leech or two and hope for the best. About 1 in 3 patients died. Once penicillin was developed and shown to be remarkably effective, correct diagnosis became important. Complaints of cough, myalgias and chills can suggest several conditions but careful auscultation revealing localized bronchial breath sounds or even something as subtle as localized crackles can point to a lung infection. It is true that these findings are not unequivocal and do require confirmatory imaging. Also, normal auscultation (like a normal history) never completely rules out a lung infection. However, virtually nothing else is as quick, easy, practically without cost and immediately available as auscultation. A rapid diagnosis of pneumonia with prompt initiation of treatment has been shown to improve outcome. Speed and availability matter!

A true story

Mr. Smith was a gentleman in his mid 50s who had a history of asthma in childhood and had begun coughing and noting shortness of breath within the past year. He went to his primary care physician who, based on the symptoms and the past history, diagnosed asthma. He began to treat him with a variety of bronchodilators and inhaled steroids and subsequently continued such treatment over the next few years. Mr. Smith's symptoms were somewhat variable but, in general, followed a downward course. Spirometry was performed and revealed a reduced vital capacity but no obstruction. In the progress notes written by his primary care physician, auscultation of the chest was always described as "clear." Finally, after

seven years of trying to treat this man's asthma without success, the physician referred him to a pulmonologist for an evaluation. At the first encounter, the pulmonologist concluded that this gentleman had far advanced pulmonary fibrosis (IPF) and referred him immediately to a transplant center. As Mr. Smith was so severely ill, he was listed quickly for transplant without having time to address his morbid obesity and deconditioning which would've been ideal. He did receive a double lung transplant. However, he suffered from multiple postoperative complications and remained in the hospital for several weeks until his death.

Mr. Smith suffered from a progressive and incurable lung disease. Although there are currently medications available that treat IPF, they are only effective in slowing the progression of the disease. By the time that Mr. Smith was eventually diagnosed, it was far too late to consider anything other than a transplant and even that was unlikely to succeed. However, had he been correctly diagnosed at an earlier stage, there would have been time to initiate one of the medications, start pulmonary rehabilitation and weight-loss regimens that might have made him a much better candidate for a lung transplant and increased his chances for several more years of life. This would have made a difference to him and to his family. His primary care physician either did not hear or did not realize the significance of inspiratory crackles that Mr. Smith must have had at the very least later in his disease course.

Another true story

A 40 year-old woman was referred to me because of a chronic cough that had persisted for two years. I have seen dozens of patients like this and determining the cause can sometimes be difficult and lengthy. During my initial examination, chest auscultation revealed profuse but faint fine crackles at both posterior bases but was otherwise unremarkable. Usually, a workup for a chronic cough starts with tests to evaluate for gastroesophageal reflux disease, Upper airway cough syndrome (UACS) or asthma.¹ Chest films or CT scans are rarely indicated. However, as crackles are unusual in this scenario, I ordered a high-resolution chest CT scan. The scan revealed diffuse reticulations and traction bronchiectasis. The pattern was not typical of idiopathic pulmonary fibrosis so the next step was a surgical biopsy that revealed chronic hypersensitivity pneumonitis (HP). HP is a rare condition but when present, often causes chronic cough. Had I not detected and recognized fine crackles and understood the implications of their presence, months might have been lost looking for common causes of a cough that were not there.

What about portable ultrasound?

ultrasound technology has been evolving rapidly and is being used in more and more situations. Some have argued that ultrasound will soon supplant the stethoscope for all physical examination screening and this will mark the end of auscultation.^{2,3} I will argue against this notion using nothing more than my brain! For purposes of this discussion, I will accept that portable ultrasound will soon rival the stethoscope in size, portability and price and that all medical students will be adequately trained in its use. What then could ultrasound detect that auscultation could not, assuming equally skilled use? Conversely, what could auscultation reveal that ultrasound could not? I've made a table that shows my personal estimates of the comparative effectiveness of auscultation vs. a hypothetical, immediately available ultrasound device in the hands of an adequately trained physician.

Psychoacoustics of auscultation

Hans Pasterkamp, MD, FRCPC (hpasterkamp@shaw.ca)

Professor, University of Manitoba, Canada

Laennec's stethoscope brought chest sounds to the attention of a much greater audience than direct auscultation had done before. His device underwent refinements over the course of the next two centuries, initially to improve its portability but later also to enhance the perception of sounds of interest, particularly in the low frequency range of heart sounds. Harvey Fletcher, who first presented the increased pressure levels required to perceive sound at low frequencies by equal loudness contours, stated that "*as machines we may describe (our ears) in the same terms that apply to devices we ourselves construct... But to understand the mechanism of the ear is by no means to understand the act of hearing, for we have not heard until the brain has perceived the message sent by the auditory nerve*" (in *Speech and Hearing*, 1929).

Psychoacoustics is a field of recognition science within psychology. Much as the study of respiratory sounds has drawn researchers from several disciplines, including physics, acoustics, electronic and computer engineering, physiology and medicine, psychoacoustics has involved these areas. However, studies of respiratory sounds that have brought researchers together, e.g. at meetings of the International Lung Sounds Association, have rarely touched on the psychoacoustics of auscultation. This review addresses relevant areas of hearing range and threshold, the characteristics of chest sounds, the masking effects of simultaneous and non-simultaneous sounds, the influence of selective hearing, and the multisensory contributions during auscultation.

Hearing range and threshold: Normal (basic) lung sounds are of low intensity during quiet breathing. Their greatest sound spectral power is in a frequency range where the perception of equal loudness requires relatively higher sound pressure levels. Respiratory effort and related changes in the sound generating air flows but also the individual anatomy as well as dynamic changes in regional ventilation and airway characteristics determine basic lung sound intensity. The perception of puerile breathing, i.e. the characteristic of basic respiratory sounds in young children and also in asthma first described by Laennec, can be explained by a relative reduction in sound spectral power at low frequencies. In asthma, this is often accompanied by an increase of sound spectral power at higher frequencies, even in the absence of wheeze. Psychologically, a decrease in lung sound intensity at low frequencies is often visualized as "reduced air entry" although this cannot explain the phenomenon in most cases of acute asthma.

Characteristics of chest sounds: Respiratory and cardiovascular sounds have the principle attributes of loudness (related to amplitude), pitch (related to frequency), duration (in relation to the respiratory phase or the cardiac cycle), and timbre (an attribute that allows to distinguish between sounds of the same perceptual loudness, pitch and duration). The latter

attribute is well recognized in cardiac auscultation, e.g. in the characterization of murmurs by their harshness, a term that is not recognized in the current formal classification of respiratory sounds. Physicians as well as caregivers of children with obstructive airway diseases are aware of "rattles, rattles and rhonchi", low pitched adventitious sounds that are often included under "wheezing" although their sound waveforms, their perception, and their relevance regarding underlying diseases and their prognoses, may be substantially different from wheezing.

Masking effects: Low intensity sounds can become inaudible in the presence of louder masking noises. An example is the increasing difficulty of recognizing brief rapidly dampened sound deflections, e.g. crackles, within random sound waves of increasing amplitude, i.e. basic lung sounds at increasing air flows. Masking effects on a sound (maskee) can also occur after an acoustic stimulus (masker). The duration of this effect depends on the duration and intensity of the masker and the relative frequency of masker and maskee. Clinically, this is of relevance during auscultation in noisy environments, including the inadvisable practice of listening through clothing.

Selective hearing: Much in our processing of acoustic signals has to do with anticipation. In speech perception, familiarity with the language is obviously of critical importance. In healthy and particularly in younger individuals, the threshold of hearing is lowest in the frequency range of speech sounds. The ability to focus on and successfully process a speech signal in a noisy environment is known as the "cocktail party effect". The intelligibility of interrupted speech depends on the interruption rate, where older and hearing-impaired subjects have most difficulties at more frequent interruptions. No studies of interobserver agreement on respiratory sounds exist from Laennec's time, but it is quite possible that he and his students understood more of the "language of the lungs" than contemporary users of the stethoscope.

Multisensory contributions: In contrast to hearing recorded audio signals of respiratory sounds, auscultation in a clinical setting provides the listener with simultaneous information from visual signals, e.g. of respiratory effort and related signs of changing respiratory resistance, of chest deformities, and of sources of potential masking noises. Basic respiratory sounds during expiration in a healthy subject at rest will be audible during only the first part of the respiratory phase while the true length of expiration until the following breath is apparent on visual inspection. Prolonged expiration in asthma is therefore, at least in part, a psychoacoustic phenomenon, based on the longer audibility of respiratory sound, particularly in the presence of wheezing.

Audio engineers are expected to have a thorough understanding of psychoacoustics. Further advances in the study and teaching of respiratory sounds will also benefit from deeper insights in this area.

Teaching and learning pulmonary auscultation – the Computerized Lung Auscultation – Sound Software (CLASS)

Alda Marques^{a,d} amarques@ua.pt; Ana Machado^{a,d}; Ana Oliveira^{a,d}; Cristina Jácome^a; Marco Pereira^{a,d}; José Moreira^b; João Rodrigues^b; José Aparício^c; Luis M. T. Jesus^{a,b}

^aLab 3R – Respiratory Research and Rehabilitation Laboratory, School of Health Sciences (ESSUA), University of Aveiro, 3810-193 Aveiro, Portugal, ^bInstitute of Electronics and Informatics Engineering of Aveiro (IEETA), University of Aveiro 3810-193 Aveiro, Portugal, ^cPaediatrics Emergency Department, Hospital Lusíadas, 4050-115 Porto, Portugal, ^dInstitute for Research in Biomedicine (iBiMED), University of Aveiro, Aveiro, Portugal

Background: Respiratory sounds (RS) are directly related to movement of air, changes within the lung tissue and position of secretions within the tracheobronchial tree, which make them valuable indicators of respiratory health. Respiratory sounds acquired through auscultation are nearly universally available, inexpensive, non-invasive and comfortable (no need to tolerate a facemask or seal around a mouthpiece), cost-effective, can be repeated as often as necessary and require minimal patient co-operation. Nevertheless, the mastering of auscultation requires complex acoustic skills to distinguish between different types of RS with similar frequencies, intensities and timings. Currently, health students are taught these skills by repeatedly listening to recordings of typical RS and visualizing their waveforms. However, these methods offer limited interaction and provide students with a narrow representativeness and usefulness of RS. Due to the subjectivity associated with the auscultation and inadequate training, the mastering of pulmonary auscultation has been decreasing and therefore, innovative teaching/learning methods to contribute for the detection/discrimination and interpretation of RS are warranted. The new era of Computer-assisted learning tools offer an exciting opportunity for self-directed learning and problem-solving, providing students with complementary activities on a computer, related with the material being taught. Such tools show great potential to be used in the teaching of pulmonary auscultation, as they would allow students to interact with a diversity of RS recorded in clinical environments, from patients with different conditions and ages, and test the knowledge acquired. However, few Computer-assisted learning tools have been developed for this purpose and they do not integrate simultaneously all the required features to enhance health students' skills on pulmonary auscultation, i.e., record, storage, playback and analysis of RS, knowledge testing and tutorials about RS.

Aim: Thus, Computerized Lung Auscultation– Sound Software (CLASS), developed for enhancing the teaching/learning of pulmonary auscultation to health professionals, will be presented.

Method: CLASS is open-source and was designed using a User-Centred Design process. This process facilitates users' workflow throughout the application as it supports their current

habits and behaviours, instead of requiring them to adapt to the application. It was developed to record, store and annotate RS in a single application and includes interactive tutorials and exercises to test the acquired knowledge. CLASS was assessed during its design process by identifying the problems that different users (medical doctors, physiotherapists, respiratory researchers and students) had when interacting with the application and the required improvements were performed throughout. Moreover, CLASS usability has been assessed by 8 physiotherapy students using computer screens recordings, think aloud reports and facial expressions according to the international standards for software validation. Time spent in each task, frequency of messages and facial expressions, number of clicks and problems reported have been counted and improvements integrated.

Results: CLASS is organized in five tabbed document interfaces: main, patients, recordings, annotations and tutorials. The “main” tab is where the information about the user and general statistics about the use of CLASS can be inserted, edited and analysed. In the “patients” tab, information about patients’ characteristics, session and recordings can be inserted, edited and analysed. The “recordings” tab allows to record (with a digital stethoscope or microphone), store and analyse RS of a selected patient and session. The “annotations” tab is where the different RS and breathing phases can be displayed, played back, annotated and analysed; and exercises, with distinct levels of complexity, can be solved based on gold standard files previously annotated by a panel of experts. Finally, the “tutorials” tab displays the definitions, acoustic properties and clinical interpretation of all types of RS.

Conclusion: CLASS is an innovative CALT and is available with all the required features for learning and consolidating pulmonary auscultation skills. It allows recording, storage, playback and analyses of RS files, practice of RS exercises, and further knowledge consolidation using the available tutorials. CLASS is simple to use and can be easily incorporated in academic activities. Moreover, since it was developed on open source components, it can be installed in individuals’ personal computers, so it can be taken to clinical or any other environments, enabling the learning process to be extended and consolidated outside academia.

Improved classification of lung sounds by medical students when spectrograms are displayed

Aviles-Solis JC (juan.c.solis@uit.no)¹, Jacome C.^{1,2}, Pasterkamp H.³, and Melbye H.¹

¹ Department of Community Medicine, UiT the Arctic University of Norway.

² Lab3R - Respiratory Research and Rehabilitation Laboratory, School of Health Sciences, University of Aveiro.

³ Dept. of Pediatrics and Child Health, University of Manitoba, Canada.

Background: Visual representation of sound with spectrograms is an attractive option to reduce variation when classifying respiratory sounds. The technology is commercially available.

Aims: To explore the impact of spectrograms on inter-observer agreement when classifying wheezes and crackles.

Methods: We used 30 recordings of respiratory sounds classified by four experienced lung sound researchers on a majority rule basis. The sample contained 15 normal recordings and 15 with wheezes or crackles. We produced spectrograms of the recordings using Adobe Audition 5.0 (Adobe Systems, San Jose, CA, USA). In a classroom, the students were shown examples of wheezes and crackles in spectrograms. Then we played the 30 recordings in a random order twice, first without the spectrogram, then with live spectrograms displayed. Twenty-three fourth year medical students at the faculty of medicine at UiT the Arctic University of Norway classified the sounds using an on-line questionnaire (Questback AS, Oslo, Norway). We asked them to classify the sounds as normal/abnormal. If abnormal, they had to further specify whether the recording contained wheezes, crackles or other sounds. We calculated kappa values for the agreement between each student and the expert classification with and without display of spectrograms, we also calculated Fleiss kappa for the 23 observers with and without the spectrograms. We used Hotelling's T^2 to explore statistically significant differences. We used the statistical software "R", version 3.2.1 and the package "multiagree".

Results: When classifying wheezes, 15/23 students (1 with $p < .05$) had a positive change in k , and 16/23 (5 with $p < .05$) when classifying crackles. All the statistically significant changes were in the direction of improved kappa values (.343 - .763). Fleiss kappa values were $k = .555$ and $k = .511$ ($p = .63$) for wheezes with and without spectrogram, respectively. For crackles, these values were $k = .403$ and $k = .223$ ($p < 0.01$) in the same order. When leaving out the five significant improvers, the Fleiss kappa values were $k = .381$ and $k = .229$ ($p = 0.01$) in the same order.

Conclusions: We found a statistically significant improvement in the agreement when classifying crackles using spectrograms. We observed no significant differences for wheezes.

Making an illustrated lung auscultation guide book for nurses: are these illustrations appropriate?

Yukio Nagasaka (rakuwadr1127@rakuwadr.com), Michiko Tsuchiya, Chikara Sakaguchi, Hitomi Ajimizu, Yosuke Nakanishi, Noboru Morikawa

Department of Pulmonary Medicine, Rakuwakai Otowa Hospital

Purpose: Recently, many textbooks and guidebooks on the auscultation of the lung are published in Japan. Most of these books do not use illustration to explain why lung sounds are generated. There are still some uncertainty why these lung sounds are generated, but simple schematic illustration will help our understanding of lung sounds. It will be needed to promote our common concepts for mechanism of lung sound generation.

Methods: We published a brief guidebook of auscultation of the lung which includes schematic illustration of lung sound generation. The illustrations in our guidebook (Figures) are shown in this presentation and will be discussed. As the real mechanisms of lung sound generation are still uncertain, the appropriateness of these illustrations should be discussed by the members of ILSA. Critical review and comments will be appreciated.

Results: Some examples of the figures are shown. Figure 1: Possible explanation of why inspiratory breath sounds are louder than expiratory breath sounds. Figure 2: Possible explanation of why frequency of wheezes in bronchial asthma fluctuates.

These figures show the possible mechanism of why these lung sounds are generated. It would be preferable if basic agreement on the illustrated mechanisms of lung sound generation will be discussed and shared by members of ILSA.

Discussion: There have been many published studies on the mechanisms of the lung sound generation. However, these results are difficult to understand for clinical personnel including physicians, nurses and physiotherapists. Simple schematic illustrations may help understanding the clinical implications of the lung sounds, and thus promote the daily use of the lung sound auscultation.

Conclusions: We propose schematic illustrations of lung sounds generation. The appropriateness of these schematic illustrations needs discussion to share our common understanding of lung sound generation.

Figures




Figure 1: Possible explanation of why inspiratory breath sounds are louder than expiratory breath sounds.

1. Inspiratory air flow hit the bifurcated bronchial wall and makes more airflow turbulence than inspiratory airflow.
2. Inspiratory airflow directs stethoscope while expiratory airflow directs opposite.




Figure 2: Possible explanation of why frequency of wheezes in bronchial asthma fluctuates.

During the expiration, the size of bronchial lumen fluctuates while the narrowest site of the bronchus (choke point) moves. These changes of the bronchial lumen result in the fluctuation of the speed of vortex shedding and frequency of wheezing.

Live Demonstration of an Advanced Pulmonary Auscultation Simulator

T.B. Talbot MD (talbot@ict.usc.edu)^{1,2}, K. Christoffersen²

¹Keck School of Medicine of the University of Southern California & ²USC Institute for Creative Technologies, Los Angeles, California, United States.

Continuing a multi-year research plan for the United States Department of Defense, we are researching the development of an advanced auscultation simulator [1] that will provide for a dynamic examination with varying acoustic output based upon listening location [2], respiratory effort & phase. The simulator will be delivered online through web browsers and support a novel pedagogical approach [3]. We employ synthetic lung & breath sounds with a clean acoustic profile so that numerous sounds can be mixed without degradation. [4].

The prototype is composed of an authoring module and a client module. We demonstrate the following capabilities of the authoring module: Respiratory loop & I:E ratio settings, selection & temporal placement of synthetic lung sound elements, layering of synthetic lung sounds, anatomic placement & acoustic listening pattern determination, and association of clinical exam elements with clinical correlation.

The client module is shown and the following are demonstrated: Clinical examination over anatomy to demonstrate changes in vesicular sounds from back, to sternum to tracheal area, various auscultation exams depicting pathologies (pneumonia, congestive heart failure, asthma, interstitial pulmonary fibrosis), clinical recognition of lung sounds with corrective feedback, and metering of exam technique for listening locations with sufficient exertion & length. Additionally, the client demonstrates use as a review tool with the ability to isolate individual sound elements.

The demonstrated design follows a cognitive task analysis of expert examinations and employs the cognitive & clinical recognition steps with a virtual mentor to provide real-time feedback to pulmonary auscultation novices.

The next stage of research will be to design and eventually conduct human research studies to determine 1) physician impressions of synthetic lung sounds for recognition and fidelity [5], 2) the assessment accuracy of the simulator versus expert assessment, and 3) the training effect of the simulator for lung sound classification [6] and clinical correlation [7].

It is hoped that these new methods for detailed assessment and individually accessible lung sound elements complexed with the proper international classification will result in improved competency & greater clinical emphasis on auscultation skills.

References

1. Talbot TB. "USC Standard Patient" accessed at www.standardpatient.org on 10 August 2017.
2. Talbot TB. A Conceptual Audio Model for Simulating Pulmonary Auscultation in a High-Fidelity Virtual Physical Examination. International Lung Sounds Association 39th Conference, Boston, 2014.

3. Talbot TB. Assessment Metrics & Performance Specifications for a Virtual Standardized Patient Comprehensive Pulmonary Auscultation Simulator. International Lung Sounds Association 40th Conference, St. Petersburg, 2015.
4. Talbot TB, Christoffersen K. Method for Live Production of Synthetic Lung Sounds in an Online Auscultation Simulator. International Lung Sounds Association 41st Conference, Tokyo, 2016.
5. Emmanouilidou D, Elhilali M. Characterization of Noise Contaminations in Lung Sound Recordings. 35th Annual International Conference of the IEEE EMBS. Osaka, 2013:2551-2554.
6. Mikami R, Murao M, Cugell DW, Chretien DW, Cole P, Meirer-Sydow J, Murphy RLH, Loudon RG. International Symposium on Lung Sounds. Chest 1987;92(2):342-345.
7. Murphy RLH. In Defense of the Stethoscope. Respiratory Care 2008;53(3):355-369.

Pulmonary Rehabilitation effects on computerized respiratory sounds of patients with AECOPD

Ana Machado^{1,2}, Ana Oliveira^{1,2}, Cátia Paixão^{1,2}, Sara Miranda^{1,2}, Hélder Melro^{1,2}, Diva Ferreira³, Alda Marques^{1,2} (amarques@ua.pt)

¹Lab3R – Respiratory Research and Rehabilitation Laboratory, School of Health Sciences, University of Aveiro, Aveiro, Portugal, ²iBiMED – Institute for Research in Biomedicine, University of Aveiro, Aveiro, Portugal,

³Pulmonology Department, Centro Hospitalar do Médio Ave, Portugal

Background: Pulmonary rehabilitation (PR) is fundamental for the management of patients with stable chronic obstructive pulmonary disease (COPD). However, its role during acute exacerbations (AECOPD) is controversial, mainly due to the lack of sensitivity of the measures used to assess changes in the respiratory system. Computerized respiratory sounds can overcome this problem as they are closely related to movement of air within the tracheobronchial tree, hence having more sensitivity to detect small changes.

Aim: To assess the effects of a community-based PR program on computerized respiratory sounds of patients with AECOPD.

Methods: 10 patients (70±8years, FEV1 46±15%predicted) with AECOPD were recruited at the emergency department and treated with standard medication plus a 3-week (2*/week) community-based PR program (i.e., breathing control and airway clearance techniques, thoracic mobility and expansion exercises, exercise training and psychoeducational support). Computerized respiratory sounds were acquired at posterior chest within 48h of the emergency episode (Pre) and following PR (Post). Adventitious respiratory sounds, i.e., wheezes and crackles, were analysed with previous validated algorithms. Differences between Pre/Post PR were explored with Wilcoxon-sign rank tests.

Results: After PR, patients presented significantly less inspiratory (Pre: 1.4 [0-5.2] vs Post: 0 [0-2.6], $p=0.016$) and expiratory crackles (Pre: 1.2 [0-21.2] vs Post: 0 [0-2], $p=0.004$), and expiratory wheezes occupation rate (Pre: 20.6 [0-82.9] vs Post: 0 [0-26.1], $p=0.004$) (Fig. 1).




Fig. 1: a) Number of crackles Pre/Post PR, per respiratory phase; b) Wheezes occupation rate Pre/Post PR, per respiratory phase.

Conclusion: Crackles and wheezes are related with increased bronchial obstruction, thus their reduction seems to indicate that PR was effective in improving the respiratory

system function. Ineffective management of AECOPD is associated with further lung function decline and higher risk of exacerbations. Therefore, PR might be an important strategy for the management of AECOPD. More studies are needed to verify these findings.

New features of 10-19 kHz sound propagation through human lungs

Vladimir Korenbaum, Anton Shiryayev, Anatoly Kostiv, Maria Safronova, Alexandr Tagiltcev, Sergei Kamenev, Alexandr Dyachenko (alexander-dyachenko@yandex.ru)
Pacific Oceanological Institute, Russian Academy of Sciences, <http://www.pio.sci.ru>

Background: The phenomenon of 10-40 kHz sound transmission through human lungs with speed of about 1000 m/s was revealed by Rueter et al., 2010. While Korenbaum et al., 2014, using signal compression technique for transmission sounding of lungs, found low- and high-speed sound propagation components in the frequency range of 10-19 kHz.

Objective: A more detailed study of the characteristics of the sound transmission of complex signals in human lungs in the frequency range of 10-19 kHz.

Method: The 14-channel apparatus (Korenbaum et al., 2014) was provided with new accelerometer sensors having a resonance frequency near 35 kHz, which made possible a linear performance in the frequency range of 10-19 kHz. Chirp signals 10-19 kHz (6 min) were emitted into human thorax (4 positions) by small shaker. A convolution of emitted and received signals technique was used. Sound propagation in human lungs was studied for opposite chest positions of shaker (ch. 1) and sensors (chs. 9-12) in 4 volunteers.

Results: The possibility of decomposition of received signals into high- and low- speed arrivals (Korenbaum et al., 2014) is verified in independent sample of subjects. The existence of low- speed arrivals with propagation velocities of 150-50 m/s, which amplitude and/or velocity is inversely dependent on an air-filling of lungs (inspiration/exhalation breath retardations) has been revealed (fig. 1a). These arrivals may be treated as the result of propagation of a sound wave through the lung parenchyma mainly. On the contrary, the amplitudes of high-speed arrivals with velocities of 150-1000 m/s are enhanced with a decrease in air-filling of the lungs during breath retardation during exhalation (fig. 1a). Thus the high-speed arrivals may be connected to the dominant sound wave propagation through high-density tissues of thorax.




Fig. 1. Transmission diagrams T_{xy} , dB on Time delay, ms – a) norm, b) reduced air-filling; blue – inspiration, green – exhalation, red dots – identified arrivals with their velocities.

Four variants of the ratio of the amplitudes of high- and low-speed arrivals and their changes depending on an air-filling of lungs (inspiration/exhalation) have been found. One of the them (fig. 1b) is characterized by the predominance of amplitudes of high-speed arrivals, both during inspiration and exhalation, which may be acoustically interpreted as a local reduction in air- filling and ventilation of lung parenchyma. It is seen only in one elderly patient with a long-term course of hormone-dependent asthma, but not in 3 other young healthy individuals.

Conclusions: The results seem to be promising for the development of high-resolution transmission acoustic imaging of lungs.

Acknowledgement: The study is supported by the RFBR grant 16-08-00075-a and the scholarship of the President of Russian Federation for graduate students and young scientists.

DIFFERENCES IN WAVEFORMS AND POWER SPECTRA OF CHEST WALL PERCUSSION SOUNDS OBTAINED BY VARIOUS SENSORS

A. Dyachenko^{1,2,3} (alexander-dyachenko@yandex.ru), M. Verenyeva^{2,3}, E. Fomina^{2,3}

¹Institute of Biomedical Problems of RAS, Moscow, Russia; ²Bauman Moscow State Technical University, Moscow, Russia; ³Prokhorov General Physics Institute of RAS, Moscow, Russia

Background: Various sensors are employed for registration of lung sounds and chest wall percussion sounds. The most widespread ones are different electronic stethoscopes with microphones. Other heavy sensors, light accelerometers and built-in microphone of a smartphone are used as well. It's important to estimate effect of sensor choice on recorded waveforms and power spectra of sounds.

Aims: To compare waveforms and power spectra of percussion tapping obtained by 1) different sensors fixed on the same place of the right back chest wall and 2) the same sensors fixed over and under diaphragm projection on the right back chest wall.

Methods: A group of 10 normal male volunteers participated in the study. Manual or apparatus taps were applied to the right back chest wall and percussion vibrations of the chest wall were registered 4.3 – 4.4 cm apart by virtue of: 1) Jabs stethoscope, 2) accelerometer, 3) one or two of 10 different smartphone models.

Results: All used sensors provided well reproducible tapping recordings. Waveforms and power spectra of tapping registered on the same place of the chest wall of the same test subject by different sensors varied considerably. Waveforms and spectra of a few pairs of sundry models of smartphones fixed over and under diaphragm projection varied much between smartphones and places of fixation. Power spectra and waveforms of accelerometer signal revealed maximums at frequencies about expected frequencies of resonance of the oscillator comprised by accelerometer mass and viscoelasticity of chest wall soft tissues.

Conclusion: Reproducibility of waveforms and spectra of registered percussion sounds is good. Differences of waveforms and spectra between sensors including various smartphones suggest standardization of sensor.

Acknowledgments: The study was supported in part by the Russian Foundation for Basic Research (RFBR) project no. 15-01-06246 and by Program IV.7.1 of the Russian Academy of Sciences (RAS) Presidium.

Using cough epochs to describe nocturnal cough in COPD patients

Patrick Fischer¹ (patrick.fischer@ges.thm.de), Lisa Steinbrecher¹, Florian Schudt¹, Keywan Sohrabi¹, Ulrich Koehler², Andreas Weissflog³, Volker Gross¹

¹ Faculty of Health Sciences, University of Applied Sciences, Giessen, ² Department of Pneumology, Intensive Care and Sleep Medicine, University Hospital of Marburg and Giessen, Marburg, Germany

³ Thora Tech GmbH, Giessen

Background and aim: Chronic cough is very common in patients suffering from chronic obstructive pulmonary disease (COPD). Several ways of quantifying cough have been published in recent years. The ERS defines a cough epoch as at least two consecutive cough events with a maximum distance of two seconds. Although in practice cough mostly occurs in epochs most of the studies focus on cough frequency quantified by counting expulsive phases. The aim of this study was to describe frequency, severity and characteristics of nocturnal cough in COPD patients, by using cough epochs.

Methods: We investigated 10 patients of each COPD stage II-IV for two consecutive nights by using objective lung sound monitoring. 11 patients reported ongoing smoking, while 19 patients quit smoking. All automatically detected cough events were manually validated and assigned to cough epochs. A classification in productive or non-productive was also done manually. Epochs containing at least one productive cough, were classified as productive.

Results: We found cough epochs in 29 of 30 patients. 75.3 ± 23.3 % of all cough events occurred in cough epochs. Most of the cough epochs were found in patients with COPD III with a median number of 6.5 productive and 8.5 normal cough epochs. Active smokers had significantly more productive cough (60.9 %) than non-smokers (24.4 %) ($p = 0.0033$). Most of the cough events were non-productive and patients showed similar distribution of cough epochs in both nights.

Conclusion: In our dataset productive cough was primarily caused by persistent smoking. With the high amount of cough events occurring in cough epochs, these cough epochs might be a promising way of quantifying cough. Especially in diseases with increased mucus production, productive cough epochs should be considered in the assessment and description of cough distribution throughout the night. This will soon be done automatically.

Investigation of pre- and postoperative lung sounds of patients who underwent general thoracic surgery

Takashi Suzuki, MD (suzuki.t@med.showa-u.ac.jp)¹, Hiroshi Kiyokawa, MD²

¹Department of General Thoracic Surgery, Showa University Hospital, ²Den-en-chofu Respiratory Clinic, Tokyo, Japan

Background: In the field of sound analysis, some low-priced software and simple voice recorders developed for music began to be available recently. Although it is difficult to calibrate these devices scientifically, they are fit to clinical use. We have auscultated, recorded, and analyzed the lung sounds of pre- and postoperative patients for 10 years.

Aims: In general thoracic surgery, it is important to find complications, for example postoperative bronchial stump fistula and retention of bronchial discharge, without delay. We have been trying to detect these disorders by analyzing the lung sounds.

Materials: The sounds of more than 100 patients undergone general thoracic surgery were analyzed.

Methods: Lung sounds were recorded in voice recorder, Linear PCM Recorder D50® [Sony, Inc. Japan], through the bilateral chest wall using a pair of hand-made stethoscopes consisting of hard gum and electret condenser microphone AT9903® [Audio-Technica, Inc. Japan]. Recorded sounds were analyzed by Adobe Audition CC®. It could easily demonstrate sound spectrogram, time-base waveform pattern, and power spectrum of the two-channel sounds.

Results: Postoperatively most of the patients, who had not have abnormal sounds preoperatively, generally did not produce adventitious sounds excluding sounds generated by mouth. Sleeve lobectomy, which consisted of resection and anastomosis of stem bronchus, produced rhonchi during both inspiration and expiration in the early postoperative course, but these rhonchi disappeared gradually. These rhonchi were supposed to be produced by retention of bronchial discharge around the anastomotic site. Postoperative bronchial fistula, which was a severe complication, generated rhonchi during expiration. These rhonchi were supposed to be produced by the gradient between intrabronchial and intrapleural pressures during expiration.

Conclusion: Lung sound analysis in general thoracic surgery provided useful information of postoperative complications.

National survey on the perception of respiratory sounds among Greek doctors (preliminary report)

Kostas Douros, Vasilis Grammeniatis, Grigorios Chatziparasidis, Antonis Platnaris, Marios Papadopoulos, [Kostas N Priftis \(MKATELA@GOLDAIR.GR\)](mailto:MKATELA@GOLDAIR.GR)

Background and aim: There are considerable differences in the perception of lung sounds among clinicians. We aimed to explore the reliance and trust in stethoscope in a sample of Greek doctors.

Methods: We constructed an electronic form that contained 5 samples of recorded adventitious lung sounds, namely, fine crackles, coarse crackles, wheezing, pleural rub, and crackles plus squawk. No clinical information or videos were given. The form was sent to 300 clinicians, mostly pediatricians, who were asked to characterize each recorded sound according to his/her own perception. Participants did not choose from predefined choices but prompted to use their own words for the description of each sound. We summarized the answers in 15 categories according to the Greek established terminology and the semantics of each term. Participants were also asked to answer how much they trusted stethoscopic findings and score their importance in the diagnostic procedure in comparison with clinical history and chest x-ray (cxr).

Results: In this first phase of the survey we received 51 completed questionnaires. Forty-eight (95.5%) reported that they trust stethoscopic findings very much. Regarding the diagnostic procedure the highest importance score (220) was given to stethoscopic findings, whereas clinical history and cxr were scored with 218 and 190 (99% and 86% of the highest score), respectively. In the recognition of adventitious sounds a correct answer was given by the majority of participants for fine and coarse crackles, and wheeze (83%, 85%, and 89%, respectively) but a variety of terms were used. Pleural rub, and the combination of crackles with squawk were recognized by 3 (5.8%) and 3 (5.8%), respectively.

Conclusion: Clinicians seem to rely on and trust stethoscope in everyday practice. Their skills in recognising adventitious sounds are quite good for the most common, but they use a confusing variety of terminology.

ISM SKRIFTSERIE - FØR UTGITT:

1. Bidrag til belysning av medisinske og sosiale forhold i Finnmark fylke, med særlig vekt på forholdene blant finskåttede i Sør-Varanger kommune.
Av Anders Forsdahl, 1976. (nytt opplag 1990)
2. Sunnhetstilstanden, hygieniske og sosiale forhold i Sør-Varanger kommune 1869-1975 belyst ved medisinalberetningene.
Av Anders Forsdahl, 1977.
3. Hjerte-karundersøkelsen i Finnmark - et eksempel på en populasjonsundersøkelse rettet mot cardiovasculære sykdommer. Beskrivelse og analyse av etterundersøkelsesgruppen.
Av Jan-Ivar Kvamme og Trond Haider, 1979.
4. D. The Tromsø Heart Study: Population studies of coronary risk factors with special emphasis on high density lipoprotein and the family occurrence of myocardial infarction.
Av Olav Helge Førde og Dag Steinar Thelle, 1979.
5. D. Reformen i distriktshelsetjenesten III: Hypertensjon i distriktshelsetjenesten.
Av Jan-Ivar Kvamme, 1980.
6. Til professor Knut Westlund på hans 60-års dag, 1983.
- 7.* Blodtrykksovervåkning og blodtrykksmåling.
Av Jan-Ivar Kvamme, Bernt Nesje og Anders Forsdahl, 1983.
- 8.* Merkesteiner i norsk medisin reist av allmennpraktikere - og enkelte utdrag av medisinalberetninger av kulturhistorisk verdi.
Av Anders Forsdahl, 1984.
9. "Balsfjordsystemet." EDB-basert journal, arkiv og statistikk-system for primærhelsetjenesten.
Av Toralf Hasvold, 1984.
10. D. Tvunget psykisk helsevern i Norge. Rettsikkerheten ved slikt helsevern med særlig vurdering av kontrollkommisjonsordningen.
Av Georg Høyer, 1986.

11. D. The use of self-administered questionnaires about food habits. Relationships with risk factors for coronary heart disease and associations between coffee drinking and mortality and cancer incidence.
Av Bjarne Koster Jacobsen, 1988.
- 12.* Helse og ulikhet. Vi trenger et handlingsprogram for Finnmark.
Av Anders Forsdahl, Atle Svendal, Aslak Syse og Dag Thelle, 1989.
13. D. Health education and self-care in dentistry - surveys and interventions.
Av Anne Johanne Sjøgaard, 1989.
14. Helsekontroller i praksis. Erfaringer fra prosjektet helsekontroller i Troms 1983-1985.
Av Harald Siem og Arild Johansen, 1989.
15. Til Anders Forsdahls 60-års dag, 1990.
16. D. Diagnosis of cancer in general practice. A study of delay problems and warning signals of cancer, with implications for public cancer information and for cancer diagnostic strategies in general practice.
Av Knut Holtedahl, 1991.
17. D. The Tromsø Survey. The family intervention study. Feasibility of using a family approach to intervention on coronary heart disease. The effect of lifestyle intervention of coronary risk factors.
Av Synnøve Fønnebø Knutsen, 1991.
18. Helhetsforståelse og kommunikasjon. Filosofi for klinikere.
Av Åge Wifstad, 1991.
19. D. Factors affecting self-evaluated general health status - and the use of professional health care services.
Av Knut Fylkesnes, 1991.
20. D. Serum gamma-glutamyltransferase: Population determinants and diagnostic characteristics in relation to intervention on risk drinkers.
Av Odd Nilssen, 1992.
21. D. The Healthy Faith. Pregnancy outcome, risk of disease, cancer morbidity and mortality in Norwegian Seventh-Day-Adventists.
Av Vinjar Fønnebø, 1992.

22. D. Aspects of breast and cervical cancer screening.
Av Inger Torhild Gram, 1992.
23. D. Population studies on dyspepsia and peptic ulcer disease: Occurrence, aetiology, and diagnosis. From The Tromsø Heart Study and The Sørreisa Gastrointestinal Disorder Studie.
Av Roar Johnsen, 1992.
24. D. Diagnosis of pneumonia in adults in general practice.
Av Hasse Melbye, 1992.
25. D. Relationship between hemodynamics and blood lipids in population surveys, and effects of n-3 fatty acids.
Av Kaare Bønnaa, 1992.
26. D. Risk factors for, and 13-year mortality from cardiovascular disease by socioeconomic status. A study of 44690 men and 17540 women, ages 40-49.
Av Hanne Thürmer, 1993.
27. Utdrag av medisinalberetninger fra Sulitjelma 1891-1990.
Av Anders Forsdahl, 1993.
28. Helse, livsstil og levekår i Finnmark. Resultater fra Hjerterundersøkelsen i 1987-88. Finnmark III.
Av Knut Westlund og Anne Johanne Sjøgaard, 1993.
29. D. Patterns and predictors of drug use. A pharmacoepidemiologic study, linking the analgesic drug prescriptions to a population health survey in Tromsø, Norway.
Av Anne Elise Eggen, 1994.
30. D. ECG in health and disease. ECG findings in relation to CHD risk factors, constitutional variables and 16-year mortality in 2990 asymptomatic Oslo men aged 40-49 years in 1972.
Av Per G. Lund-Larsen, 1994.
31. D. Arrhythmia, electrocardiographic signs, and physical activity in relation to coronary heart risk factors and disease. The Tromsø Study.
Av Maja-Lisa Løchen, 1995.
32. D. The Military service: mental distress and changes in health behaviours among Norwegian army conscript.
Av Edvin Schei, 1995.

33. D. The Harstad injury prevention study: Hospital-based injury recording and community-based intervention.
Av Børge Ytterstad, 1995.
- 34.* D. Vilkår for begrepsdannelse og praksis i psykiatri. En filosofisk undersøkelse.
Av Åge Wifstad, 1996.
(utgitt Tano Aschehoug forlag 1997)
35. Dialog og refleksjon. Festskrift til professor Tom Andersen på hans 60-års dag, 1996.
36. D. Factors affecting doctors' decision making.
Av Ivar Sønbo Kristiansen, 1996.
37. D. The Sørreisa gastrointestinal disorder study. Dyspepsia, peptic ulcer and endoscopic findings in a population.
Av Bjørn Bernersen, 1996.
38. D. Headache and neck or shoulder pain. An analysis of musculoskeletal problems in three comprehensive population studies in Northern Norway.
Av Toralf Hasvold, 1996.
39. Senfølger av kjernefysiske prøvespreninger på øygruppen Novaya Semlya i perioden 1955 til 1962. Rapport etter programmet "Liv". Arkangelsk 1994.
Av A.V. Tkatchev, L.K. Dobrodeeva, A.I. Isaev, T.S. Podjakova, 1996.
40. Helse og livskvalitet på 78 grader nord. Rapport fra en befolkningsstudie på Svalbard høsten 1988.
Av Helge Schirmer, Georg Høyer, Odd Nilssen, Tormod Brenn og Siri Steine, 1997.
- 41.* D. Physical activity and risk of cancer. A population based cohort study including prostate, testicular, colorectal, lung and breast cancer.
Av Inger Thune, 1997.
42. The Norwegian - Russian Health Study 1994/95. A cross-sectional study of pollution and health in the border area.
Av Tone Smith-Sivertsen, Valeri Tchachtchine, Eiliv Lund, Tor Norseth, Vladimir Bykov, 1997.
43. D. Use of alternative medicine by Norwegian cancer patients
Av Terje Risberg, 1998.

43. D. Incidence of and risk factors for myocardial infarction, stroke, and diabetes mellitus in a general population. The Finnmark Study 1974-1989.
Av Inger Njølstad, 1998.
45. D. General practitioner hospitals: Use and usefulness. A study from Finnmark County in North Norway.
Av Ivar Aaraas, 1998.
- 45B Sykestuer i Finnmark. En studie av bruk og nytteverdi.
Av Ivar Aaraas, 1998.
46. D. No går det på helsa laus. Helse, sykdom og risiko for sykdom i to nord-norske kystsamfunn.
Av Jorid Andersen, 1998.
47. D. The Tromsø Study: Risk factors for non-vertebral fractures in a middle-aged population.
Av Ragnar Martin Joakimsen, 1999.
48. D. The potential for reducing inappropriate hospital admissions: A study of health benefits and costs in a department of internal medicine.
Av Bjørn Odvar Eriksen, 1999.
49. D. Echocardiographic screening in a general population. Normal distribution of echocardiographic measurements and their relation to cardiovascular risk factors and disease. The Tromsø Study.
Av Henrik Schirmer, 2000.
50. D. Environmental and occupational exposure, life-style factors and pregnancy outcome in arctic and subarctic populations of Norway and Russia.
Av Jon Øyvind Odland, 2000.
- 50B Russisk utgave av Skriftserie 50
51. D. A population based study on coronary heart disease in families. The Finnmark Study 1974-1989.
Av Tormod Brenn, 2000.
52. D. Ultrasound assessed carotid atherosclerosis in a general population. The Tromsø Study.
Av Oddmund Joakimsen, 2000.

53. D. Risk factors for carotid intima-media thickness in a general population. The Tromsø Study 1979-1994.
Av Eva Stensland-Bugge, 2000.
54. D. The South Asian cataract management study.
Av Torkel Snellingen, 2000.
55. D. Air pollution and health in the Norwegian-Russian border area.
Av Tone Smith-Sivertsen, 2000.
56. D. Interpretation of forearm bone mineral density. The Tromsø Study.
Av Gro K. Rosvold Berntsen, 2000.
57. D. Individual fatty acids and cardiovascular risk factors.
Av Sameline Grimsgaard, 2001.
58. Finnmarkundersøkelsene
Av Anders Forsdahl, Fylkesnes K, Hermansen R, Lund E, Lupton B, Selmer R, Straume E, 2001.
59. D. Dietary data in the Norwegian women and cancer study. Validation and analyses of health related aspects.
Av Anette Hjartåker, 2001.
60. D. The stenotic carotid artery plaque. Prevalence, risk factors and relations to clinical disease. The Tromsø Study.
Av Ellisiv B. Mathiesen, 2001.
61. D. Studies in perinatal care from a sparsely populated area.
Av Jan Holt, 2001.
62. D. Fragile bones in patients with stroke? Bone mineral density in acute stroke patients and changes during one year of follow up.
Av Lone Jørgensen, 2001.
63. D. Psychiatric morbidity and mortality in northern Norway in the era of deinstitutionalisation. A psychiatric case register study.
Av Vidje Hansen, 2001.
64. D. Ill health in two contrasting countries.
Av Tom Andersen, 1978/2002.
65. D. Longitudinal analyses of cardiovascular risk factors.
Av Tom Wilsgaard, 2002.

66. Helseundersøkelsen i Arkhangelsk 2000.
Av Odd Nilssen, Alexei Kalinin, Tormod Brenn, Maria Averina et al., 2003.
67. D. Bio-psycho-social aspects of severe multiple trauma.
Av Audny G. W. Anke, 2003.
68. D. Persistent organic pollutants in human plasma from inhabitants of the arctic.
Av Torkjel Manning Sandanger, 2003.
69. D. Aspects of women's health in relation to use of hormonal contraceptives and pattern of child bearing.
Av Merethe Kumle, 2003.
70. Pasienterfaringer i primærlegetjenesten før og etter fastlegereformen.
Av Olaug Lian, 2003.
71. D. Vitamin D security in northern Norway in relation to marine food traditions.
Av Magritt Brustad, 2004.
72. D. Intervensjonsstudien i Finnmark. Evaluering av lokalsamfunns basert hjerte- og kar forebygging i kystkommunene Båtsfjord og Nordkapp.
Av Beate Lupton, 2004.
73. D. Environmental factors, metabolic profile, hormones and breast and endometrial cancer risk.
Av Anne-Sofie Furberg, 2004.
74. D. Det skapende mellomrommet i møtet mellom pasient og lege.
Av Eli Berg, 2004.
75. Krefregisteret i Arkhangelsk oblast i nordvest Russland. Med en sammenligning av kreftforekomst i Arkhangelsk oblast og Norge 1993 - 2001.
Av Vaktskjold Arild, Lebedintseva Jelena, Korotov Dmitrij, Tkatsjov Anatolij, Podjakova Tatjana, Lund Eiliv, 2004
76. D. Characteristics and prognosis of long-term stroke survivors. The Tromsø Study.
Av Torgeir Engstad, 2004
77. D. Withdrawal and exclusion. A study of the spoken word as means of understanding schizophrenic patients.
Av Geir Fagerjord Lorem, 2005.

78. "Søkelys på samfunnsmedisinene." Evaluering av kommunal samfunnsmedisinsk legetjeneste, offentlig legearbeid og de forebyggende oppgaver i Fastlegeordningen.
Av Betty Pettersen og Roar Johnsen, 2005.
79. Prosjekt egenmelding Kristiansand kommune. Evaluering av kontrollert intervensjonsforsøk i stor skala, med utvidet rett til egenmelding i kombinasjon med økt og formalisert samhandling mellom arbeidstaker og arbeidsplassen ved sykefravær.
Av Nils Fleten og Roar Johnsen, 2005.
80. D. Abdominal aortic aneurysms: Diagnosis and epidemiology. The Tromsø study.
Av Kulbir Singh, 2005.
81. D. A population based study on cardiovascular diseases in Northwest Russia. The Arkhangelsk study 2000.
Av Maria Averina, 2005.
82. D. Exposure to exogenous hormones in women: risk factors for breast cancer and molecular signature.
Av Vanessa Dumeaux, 2005.
83. D. Repeated ultrasound measurements of carotid artery plaques in a general population. The Tromsø Study 1994-2001.
Av Stein Harald Johnsen, 2005.
84. D. Risk Factors For Fractures In Tromsø. The Tromsø Study.
Av Luai Awad Ahmed, 2005.
85. D. The quality and use of two health registries in Russia. The Arkhangelsk Cancer Registry and the Kola Birth Registry
Качество и использование двух медицинских регистров в России. Архангельск регистр рака и Кольский регистр родов
Av Arild Vaktskjold, 2005.

86. D. Haemoglobin, anaemia and haematological malignancies.
Av Tove Skjelbakken, 2006
87. D. The sick-listed - an under-recognised resource in handling sickness absence.
Av Nils Fleten, 2006.
88. D. Longitudinal changes in forearm bone mineral density in women and men from 25 to 84 years.
The Tromsø Study.
Av Nina Emaus, 2006.
89. D. Asthma and allergy in children. An epidemiological study of asthma and allergy in schoolchildren living in Northern Norway and Russia with respect to prevalence trends 1985-1995-2000, geographic differences in prevalence and biomarkers.
By Anders Selnes, 2006.
90. D. "Nå ska du høre ka æ mene med arv." Samisk forståelse av arv som en utfordring i medisinsk genetikk.
Av Valeria Marton, 2006 -
Senter for Samisk Helseforskning
91. D. Sex steroids, bone loss and non-vertebral fractures in women and men. The Tromsø Study.
By Åshild Bjørnerem, 2007.
92. D. Substance use behaviour among ethnic diverse young people in North Norway in the 1990s.
"The North Norwegian Youth Study": A cross-cultural longitudinal study comparing smoking and drinking rates and patterns among young indigenous Sami and non-indigenous peers
Av Anna Rita Spein, 2007.
Senter for Samisk Helseforskning
93. D. Infection, inflammation and atherosclerosis.
Av Dag S. Halvorsen, 2007.
94. D. Hormones, Smoking and Mammographic Density in Postmenopausal Norwegian Women.
The Tromsø Mammography and Breast Cancer Study.
Av Yngve Bremnes, 2007.
95. D. Suicidal behaviour among indigenous Sami in Arctic Norway. A special focus on adolescents and young adults.
Av Anne Silviken, 2007.

96. D. Explaining the socioeconomic variation in incidence and survival of cancer. Analyses and multiple imputation of data from The Norwegian Women and Cancer Study and The Norwegian-Swedish Women's Lifestyle and Health Cohort Study.
Av Tonje Braaten, 2008.
97. D. Local Public Health Physicians in Norway from 1994 to 2002. Workload, work content, and interaction. A story of everyday life in primary health care.
Av Betty Johanne Pettersen, 2008.
98. D. Iron status and prevalence of hereditary haemochromatosis in a multiethnic population in northern Norway. The SAMINOR study, The Sør-Varanger study, The Tromsø V study
Av Ann Ragnhild Broderstad, 2008.
99. D. The consumption of lean and fatty fish, different dietary patterns, and the risk of cancers of various sites.
Av Dagrun Engeset. 2008.
100. D. Coercion in the delivery of mental health services in Norway.
Av Knut Ivar Iversen, 2008.
101. DD. Explaining risk reductions in medical practice: Prevention or postponement?
Av Peder Andreas Halvorsen, 2008.
102. D. Ankylosing spondylitis, aortic regurgitation, acetabular dysplasia and osteoarthritis of the hip. An epidemiological survey in a Norwegian Sámi population.
Av Knut Johnsen, 2009.
103. D. Helicobacter pylori and dyspepsia from a public health perspective. The Sørreisa gastrointestinal disorder study.
Av Anne Mette Asfeldt, 2009.
104. D. The Murmansk County Birth Registry (MCBR) The implementation and applicability of a population-based medical birth registry in the Russian Arctic
Av Erik Eik Anda, 2009.

105. D. Diet, dietary supplements and dietary change in cancer survivors and cancer-free persons - the Norwegian Women and Cancer study and the European Prospective Investigation into Cancer and Nutrition
Av Guri Skeie, 2009.
106. D. Some issues of provision and access to dental services in Norway
Av Birgit Abelsen, 2009.
107. D. A population-based study on cardiovascular risk factors and self-reported type 2 diabetes mellitus in the sami population. The SAMINOR Study.
Av Tove Anita Nystad, 2010.
108. D. Human exposure to perfluorinated compounds concentrations, dietary impact and molecular signatures.
Av Charlotta Rylander, 2010.
109. D. Hormone therapy use, sex hormone concentrations and gene expression - The Norwegian Women and Cancer study (NOWAC)
Av Marit Waaseth, 2010.
110. D. Abdominal aortic aneurysms: Risk factors and cardiovascular consequences. The Tromsø Study.
Av Signe Helene Forsdahl, 2010.
111. D. Young Adults and Seafood. Using the voice of consumers to develop new seafood product concepts aimed at increasing consumption
Av Themistoklis Altintzoglou, 2010.
112. D. The development and use of a new tool for estimating individual sun included vitamin D in epidemiological surveys.
Av Kåre Edvardsen, 2010.
113. D. Incidence of and risk factors for type 2 diabetes in a general population. The Tromsø Study.
Av Josepha Joseph, 2011.
114. D. Childhood Abuse - Pregnancy and Childbirth.
Av Mirjam Lukasse, 2011.
115. D. Ethnic discrimination and bullying in relation to self-reported physical and mental health in Sami settlement areas in Norway. The Saminor study
Ketil Lenert Hansen, 2011.

116. D. Body mass index and risk of malnutrition in community-living elderly men and women: Relationships with morbidity, mortality and health-related quality of life. The Tromsø and HUNT studies.
Av Jan-Magnus Kvamme, 2011.
117. D. Factors behind high cardiovascular disease mortality in Northwest Russia The Arkhangelsk study
Av Oleg Sidorenkov, 2011.
118. D. Functional ultrasound of the anal canal. The effect of pregnancy and childbirth.
Av Ingrid Petrikke Olsen, 2011.
119. D. Maternal concentrations, predictors and change in profiles of organochlorines, toxic and essential elements during pregnancy and postpartum. The Vietnamese Mother-and-Child Study and the Northern Norwegian Mother-and-Child Study.
Av Solrunn Hansen, 2011.
120. D. Morally bound medical work. An empirical study exploring moral conditions of doctors' everyday practice.
Av Kari Milch Agledahl, 2011.
121. D. A community-based factorial trial on Alzheimer's disease. Effects of expectancy, recruitment methods, co-morbidity and drug use. The Dementia Study in Northern Norway.
Av Fred Andersen, 2011.
122. D. Physical activity, osteoporosis, and fracture risk. Long-term associations in a general population.
Av Bente Morseth, 2011.
123. DD. Why don't we take a look at the patient? An anthropological analysis of how doctors become doctors.
Av Torstein Risør, 2011.
124. D. Aspects of health services in Sami areas.
Av Margrete Gaski, 2012.
125. D. Disease activity and outcome in juvenile idiopathic arthritis; a longitudinal cohort study in the Nordic countries.
Av Ellen Berit Nordal, 2012.

126. D. Carotid atherosclerosis, vascular risk factors and relation to cognitive test results. The Tromsø Study 1994-2008.
Av Kjell Arne Arntzen, 2012.
127. D. A doctor close at hand. A qualitative analysis of GPs' work in cancer care.
Av May-Lill Johansen, 2012.
128. D. Seasonal variation in incidence of acute myocardial infarction and cardiovascular disease risk factors in a subarctic population. The Tromsø Study.
Av Laila Arnesdatter Hopstock, 2012.
129. D. Nutrients and toxic elements in semi-domesticated reindeer in Norway. Nutritional and food safety aspects.
Av Amar Eltayeb Ali Hassan, 2012.
130. D. COPD in the elderly - diagnostic criteria, symptoms and smoking. Quantitative and qualitative studies of persons 60 years and older in The Tromsø studies.
Av Astri Medbø, 2012.
131. D. A study of fish consumption and cardiometabolic risk factors among the circumpolar population of the rural Nenets Autonomous Area in comparison with the urban population of Arkhangelsk County.
Av Natalia Petrenya, 2012.
132. D. Road traffic crashes in Arkhangelsk, Russia in 2005-2010.
Av Alexander Kudryavtsev, 2013.
133. D. Respiratory symptoms, lung function, and occupational exposure among seafood industry workers
A study among employees at Norwegian salmon factories and Russian North-West trawl fleet (Arkhangelsk)
Av Olga Shiryaeva, 2013.
134. D. Genetic susceptibility, lifestyle, ovarian hormones, and mammographic density - key factors in breast cancer development. The Norwegian Energy Balance and Breast Cancer Aspects-I Study.
Av Anita Iversen, 2013.

135. D. Physical activity, mortality and breast cancer risk.
The Norwegian Women and Cancer Study.
Av Kristin Benjaminsen Borch, 2013.
136. D. Retinopathy with and without diabetes: Risk factors and visual impairment.
The Tromsø Eye Study and a Norwegian screening study.
Av Geir Bertelsen, 2013.
137. D. Age-related macular degeneration: Prevalence and risk factors - a cross-sectional study. The Tromsø Study 2007/2008.
Av Maja Gran Erke, 2013.
138. D. Staphylococcus aureus nasal carriage - Interplay between host, microbe and the environment.
- The Tromsø Staph and Skin Study.
Av Karina Olsen, 2013.
139. D. Blood gene expression, lifestyle and diet. The Norwegian Women and Cancer Post-genome Cohort.
Av Karina Standahl Olsen, 2013.
140. D. Human biomonitoring of perfluoroalkyl substances and cyclic volatile methylsiloxanes. Concentrations in plasma, serum and whole blood from pregnant, delivering or postmenopausal women, and cord blood.
Av Linda Hanssen, 2013.
141. D. A population-based study of health care utilisation according to care level, socio-economic group, and continuity of primary care. The Tromsø Study.
Av Anne Helen Hansen, 2013.
142. D. Social determinants of self-rated health and cardiovascular disease among the Sami and other Arctic indigenous peoples.
The SLiCA study and the SAMINOR study
Av Bent-Martin Eliassen, 2013.
143. D. Vertebral fractures: prevalence, risk factors, and health-related quality of life.
The Tromsø Study 2007/2008.
Av Svanhild Waterloo, 2013.

144. D. Genital *chlamydia trachomatis* infections among adolescents in a high-incidence area in Norway: genotypes, prevalence, early sexual behavior and testing patterns - a cross-sectional study. The Finnmark High School Study (FHSS)
Av Kirsten Gravningen, 2013.
145. D. Associations between Primary Health Care - and Hospital Utilization among elderly People in Norway.
Av Trygve Sigvart Deraas, 2013.
146. D. Prenatal exposure to DDT and other selected environmental contaminants and their predictors in malaria and non-malaria areas in coastal KwaZulu Natal, South Africa.
Av Kalavati Channa, 2014.
147. D. Predictors of progression of ultrasound-assessed carotid artery atherosclerosis. The Tromsø Study 1994-2008
Av Marit Herder, 2014.
148. D. A study of occupational health among mine workers in Murmansk Oblast, Russia.
Av Morten Skandfer, 2014.
149. D. Psoriasis, overweight and metabolic syndrome. The Tromsø Study.
Av Kjersti Danielsen, 2014.
150. D. Smoking and incidence and mortality of colorectal cancer.
Av Ranjan Parajuli, 2014.
151. D. Understanding temporality in human concentrations of organic contaminants. Considering human concentrations over time and through life in perspective of historic production and use.
Av Therese Haugdahl Nøst, 2014.
152. D. The Clinician in Leadership. Perceptions of Style. Perspectives from Rural Primary Medicine in Norway.
Av Jan Hana, 2014.
153. D. Aspects of Active Smoking and Breast Cancer.
Av Eivind Bjerkaas, 2014.

154. D. Fractional exhaled nitric oxide and its relation to exercise, asthma and allergic rhinoconjunctivitis in a subarctic childhood population.
A study of asthma and allergy among schoolchildren in Nordland County
Av Bjørg Evjenth, 2014.
155. D. Managing childhood obesity
The Finnmark Activity School.
Av Ane Sofie Kokkvoll, 2014.
156. D. "Light my fire" - perspectives on motivation, helpfulness and implementation of a guided internet-based cognitive behavioral therapy.
Av Maja Wilhelmsen, 2015.
157. D. Pulse oximetry in an adult population: Predictors of low oxygen saturation and associations with mortality.
The Tromsø Study 2001-2008.
Av Monica Linea Vold, 2015.
158. D. General Practitioners' Decisions to Refer Patients to Secondary Care - Referral Rates, Reasons for Referral and Expected Medical Benefit of the Referrals
Av Unni Ringberg, 2015.
159. D. Persistent post-surgical pain
Prevalence, risk factors and pain mechanisms
Av Aslak Johansen, 2015.
160. D. Remuneration and organization in general practice: Three essays on doctors' preferences
Av Jon Helgheim Holte, 2015.
161. D. Sámi ethnicity as a variable.
Premises and implications for population-based studies on health and living conditions in Norway
Av Torunn Pettersen, 2015.
162. D. Concentrations and predictors of persistent organic pollutants in pregnant women and associations with maternal and infant thyroid homeostasis.
The Northern Norway Mother-and-Child Contaminant Cohort Study.
Av Vivian Berg, 2015.

163. D. Atrial fibrillation in The Tromsø Study 1994-2007. Risk factors, occurrence and gender differences.
Av Audhild Nyrnes, 2016.
164. D. Comparisons of sickness absence patterns, trends and attitudes in the health and care sectors in two municipalities in Norway and Denmark.
Av Line Krane, 2016.
165. D. The Northern Norway Mother-and-Child Contaminant Cohort study (MISA): Population Characteristics, Dietary Intake, and Predictors of Organochlorine Contaminants in Meconium and Maternal Serum and of Essential and Toxic Elements in Mothers' Whole Blood.
Av Anna Sofia Veyhe, 2016.
166. Du mao berre putla pao. Olav Helge Førde - 70 år. Ei festminnebok.
167. D. Use of analgesics in the general population. Trends, persistence, high-risk use and associations with pain sensitivity
Av Per-Jostein Samuelsen, 2016.
168. D. Potato consumption and risk of colorectal and pancreatic cancer. The Norwegian Women and Cancer cohort and the HELGA cohort.
Av Lene Angell Åsli, 2016.
169. D. Suicides in The Nenets Autonomous Okrug, Russia.
Av Yuri Alexandrovich Sumarokov, 2016.
170. D. Experiences with outpatient commitment orders from the perspectives of patients, relatives and staff. A qualitative study.
Av Bjørn Stensrud, 2016.
171. D. Using medical birth registries in the Kola Peninsula for birth defects surveillance and investigation of their risk factors
Av Vitaly Postoev, 2016.
172. D. Whole-grain bread, milk, and risk of colorectal cancer. The Norwegian Women and Cancer Study.
Av Toril Bakken, 2016.
173. Psykomotorisk fysioterapi - en praksis i bevegelse. Festskrift til Gudrun Øvreberg.

174. D. When coercion moves into your home.
A study of outpatient commitment in Northern Norway
Av Henriette Riley, 2016.
175. D. Training interactions in local teams:
Using critical participatory action research
to explore context based learning
Av Helen Brandstorp, 2017.
176. D. Adolescents and self-harm.
A study of factors associated with suicide and use of
health care services following self-harm in national
representative populations of Norwegian adolescents
Av Elin Anita Fadum, 2017.
177. D. Decline in resting heart rate, its association with
other variables, and its role in cardiovascular
disease.
The Tromsø Study
Av Ekaterina Sharashova, 2017.
178. D. Risk factors of adverse pregnancy outcomes:
opportunities and perspectives of a birth registry-
based study
Av Anna Usynina, 2017.
179. D. Managing exacerbations of chronic obstructive
pulmonary disease.
An investigation of treatment decisions from a
primary care perspective.
Av Johanna Laue, 2017.
180. D. "Breaking the silence"
Interpersonal violence and Health among Sami and non-
Sami. A population-based study in Mid- and Northern
Norway.
Av Astrid M.A. Eriksen, 2017.
181. D. self-reported musculoskeletal complaints. Prevalence,
risk factors, and mortality.
Av Ole Fredrik Linnemann Andorsen, 2017.

De som er merket med D er doktorgradsarbeid.
De som er merket med DD er doktorgradsarbeid fra universitet
i utlandet. De som er merket med * har vi dessverre ikke flere
eksemplarer av.